



BAAP OF ALL FORMULA LISTS






FOR IIT JEE

INTEGRAL CALCULUS

[Download Doubtnut Today](#)

SL#	FORMULA
1	$\int f(x)dx = F(x) + C$ if $F'(x) = f(x)$.
2	$\left(\int f(x)dx\right)' = f(x)$
3	$\int kf(x)dx = k\int f(x)dx$
4	$\int [f(x) + g(x)]dx = \int f(x)dx + \int g(x)dx$
5	$\int [f(x) - g(x)]dx = \int f(x)dx - \int g(x)dx$
6	$\int f(ax)dx = \frac{1}{a}F(ax) + C$
7	$\int f(ax + b)dx = \frac{1}{a}F(ax + b) + C$
	DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE
8	$\int f(x)df'(x)dx = \frac{1}{2}f^2(x) + C$
9	$\int \frac{f'(x)}{f(x)}dx = \ln f(x) + C$
10	Method of Substitution $\int f(x)dx = \int f(u(t))u'(t)dt$ if $x = u(t)$.
11	Integration by Parts $\int u dv = uv - \int v du$, where $u(x), v(x)$ are differentiable functions.
12	$\int adx = ax + C$
13	$\int x dx = \frac{x^2}{2} + C$
14	$\int x^2 dx = \frac{x^3}{3} + C$
15	$\int x^p dx = \frac{x^{p+1}}{p+1} + C, p \neq -1$
	DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE
16	$\int (ax + b)^n dx = \frac{(ax + b)^{n+1}}{a(n+1)} + C, n \neq -1$.
17	$\int \frac{dx}{x} = \ln x + C$
18	$\int \frac{dx}{ax + b} = \frac{1}{a}\ln ax + b + C$

19	$\int \frac{ax+b}{cx+d} dx = \frac{a}{c}x + (bc-ad)c^2 \ln cx+d + C$
20	$\int \frac{dx}{x+a}(x+b) = \frac{1}{a-b} \ln \left \frac{x+b}{x+a} \right + C, a \neq b.$
21	$\int \frac{xdx}{a+bx} = \frac{1}{b^2}(a+bx - \ln a+bx) + C$
22	$\int \frac{x^2 dx}{a+bx} = \frac{1}{b^3} \left[\frac{1}{2}(a+bx)^2 + a^2 \ln a+bx \right] + C$
23	$\int \frac{dx}{x(a+bx)} = \frac{1}{a} \ln \left \frac{a+bx}{x} \right + C$
 पढ़ना हुआ आसान	📄 DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE
24	$\int \frac{dx}{x^2(a+bx)} = -\frac{1}{ax} + \frac{b}{a^2} \ln \left \frac{a+bx}{x} \right + C$
25	$\int \frac{xdx}{a+bx} = \frac{1}{b^2} \left(\ln a+bx + \frac{a}{a+bx} \right) + C$
26	$\int \frac{x^2 dx}{a+bx^2} = \frac{1}{b^3} \left(a+bx - 2a \ln a+bx - \frac{a^2}{a+bx} \right) + C$
27	$\int \frac{dx}{x(a+bx)^2} = \frac{1}{a(a+bx)} + \frac{1}{a^2} \ln \left \frac{a+bx}{x} \right + C$
28	$\int \frac{dx}{x^2-1} = \frac{1}{2} \ln \left \frac{x-1}{x+1} \right + C$
29	$\int \frac{dx}{1-x^2} = \frac{1}{2} \ln \left \frac{1+x}{1-x} \right + C$
30	$\int \frac{dx}{a^2-x^2} = \frac{1}{2a} \ln \left \frac{a+x}{a-x} \right + C$
31	$\int \frac{dx}{x^2-a^2} = \frac{1}{2a} \ln \left \frac{x-a}{x+a} \right + C$
 पढ़ना हुआ आसान	📄 DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE
32	$\int \frac{dx}{1+x^2} = \tan^{-1} x + C$
33	$\int \frac{dx}{a^2+x^2} = \frac{1}{a} \frac{\tan^{-1} x}{a} + C$
34	$\int \frac{xdx}{x^2+a^2} = \frac{1}{2} \ln(x^2+a^2) + C$
35	$\int \frac{dx}{a+bx^2} = \frac{1}{\sqrt{ab}} \tan^{-1} \left(x \sqrt{\frac{b}{a}} \right) + C, ab > 0$
36	$\int \frac{xdx}{a+bx^2} = \frac{1}{2b} \ln \left x^2 + \frac{a}{b} \right + C$
37	$\int \frac{dx}{x(a+bx^2)} = \frac{1}{2a} \ln \left \frac{x^2}{a+bx^2} \right + C$
38	$\int \frac{dx}{a^2-b^2x^2} = \frac{1}{2ab} \ln \left \frac{a+bx}{a-bx} \right + C$
39	$\int \frac{dx}{ax^2+bx+c} = \frac{1}{\sqrt{b^2-4ac}} \ln \left \frac{2ax+b-\sqrt{b^2-4ac}}{2ax+b+\sqrt{b^2-4ac}} \right + C, b^2-4ac > 0$
 पढ़ना हुआ आसान	📄 DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE
40	$\int \frac{dx}{ax^2+bx+c} = \frac{2}{\sqrt{4ac-b^2}} \frac{\tan^{-1}(2ax+b)}{\sqrt{4ac-b^2}} + C, b^2-4ac < 0$

41	$\int \frac{dx}{\sqrt{ax+b}} = \frac{2}{a}\sqrt{ax+b} + C$
42	$\int \sqrt{ax+bdx} = \frac{2}{3a}(ax+b)^{\frac{3}{2}} + C$
43	$\int \frac{xdx}{\sqrt{ax+b}} = \frac{2(ax-2b)}{3a^2}\sqrt{ax+b} + C$
44	$\int x\sqrt{ax+bdx} = \frac{2(3ax-2b)}{15a^2}(ax+b)^{\frac{3}{2}} + C$
45	$\int \frac{dx}{(x+c)\sqrt{ax+b}} = \frac{1}{\sqrt{b-ac}} \ln \left \frac{\sqrt{ax+b} - \sqrt{b-ac}}{\sqrt{ax+b} + \sqrt{b-ac}} \right + C, b-ac > 0.$
46	$\int \frac{dx}{(x+c)\sqrt{ax+b}} = \frac{1}{\sqrt{ac-b}} \tan^{-1} \sqrt{\frac{ax+b}{ac-b}} + C, b-ac < 0.$
47	$\int x^2 \sqrt{a+bx} dx = \frac{2(8a^2 - 12abx + 15b^2x^2)}{105b^3} \sqrt{(a+bx)^3} + C$



[DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE](#)

48	$\int \frac{x^2 dx}{\sqrt{a+bx}} = \frac{2(8a^2 - 4abx + 3b^2x^2)}{15b^3} \sqrt{a+bx} + C$
49	$\int \frac{dx}{x\sqrt{a+bx}} = \frac{1}{\sqrt{a}} \ln \left \frac{\sqrt{a+bx} - \sqrt{a}}{\sqrt{a+bx} + \sqrt{a}} \right + C, a > 0$
50	$\int \frac{dx}{x\sqrt{a+bx}} = \frac{2}{\sqrt{-a}} \tan^{-1} \left \frac{a+bx}{-a} \right + C, a < 0.$
51	$\int \sqrt{\frac{a-x}{b+x}} dx = \sqrt{(a-x)(b+x)} + (a+b) \sin^{-1} \sqrt{\frac{x+b}{a+b}} + C$
52	$\int \sqrt{\frac{a+x}{b-x}} dx = -\sqrt{(a+x)(b-x)} - (a+b) \sin^{-1} \sqrt{\frac{b-x}{a+b}} + C$
53	$\int \frac{dx}{\sqrt{x-a}(b-a)} = 2 \sin^{-1} \sqrt{\frac{x-a}{b-a}} + C$
54	$\int \sqrt{a+bx-cx^2} dx = \frac{2cx-b}{4c} \sqrt{a+bx-cx^2} + \frac{b^2-4ax}{8\sqrt{c^3}} \frac{\sin^{-1}(2cx-b)}{\sqrt{b^2+4ac}} + C$
55	$\int \frac{dx}{\sqrt{ax^2+bx+c}} = \frac{1}{\sqrt{a}} \ln 2ax+b+2\sqrt{ax^2+bx+c} + C, a > 0$



[DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE](#)

56	$\int \frac{dx}{\sqrt{ax^2+bx+c}} = -\frac{1}{\sqrt{a}} \frac{\sin^{-1}(2ax+b)}{4a} \sqrt{b^2-4ac} + C, a < 0$
57	$\int \sqrt{x^2+a^2} dx = \frac{x}{2} \sqrt{x^2+a^2} + \frac{a^2}{2} \ln x + \sqrt{x^2+a^2} + C$
58	$x\sqrt{x^2+a^2} dx = \frac{1}{3} (x^2+a^2)^{\frac{3}{2}} + C$
59	$\int x^2 \sqrt{x^2+a^2} dx = \frac{x}{8} (2x^2+a^2) \sqrt{x^2+a^2} - \frac{a^4}{8} \ln x + \sqrt{x^2+a^2} + C$
60	$\int \frac{\sqrt{x^2+a^2}}{x^2} dx = -\frac{\sqrt{x^2+a^2}}{x} + \ln x + \sqrt{x^2+a^2} + C$
61	$\int \left(\frac{\sqrt{x^2+a^2}}{x} \right) dx = \sqrt{x^2+a^2} + a \ln \left \frac{x}{a + \sqrt{x^2+a^2}} \right + C$
62	$\int \frac{xdx}{\sqrt{x^2+a^2}} = \sqrt{x^2+a^2} + C$
63	

$$\int \frac{x^2 dx}{\sqrt{x^2 + a^2}} = \frac{x}{2} \sqrt{x^2 + a^2} - \frac{a^2}{2} \ln|x + \sqrt{x^2 + a^2}| + C$$

64
$$\int \frac{dx}{x\sqrt{x^2 + a^2}} = \frac{1}{a} \ln \left| \frac{x}{a + \sqrt{x^2 + a^2}} \right| + C$$

65
$$\int \sqrt{x^2 - a^2} dx = \frac{x}{2} \sqrt{x^2 - a^2} - \frac{a^2}{2} \ln|x + \sqrt{x^2 - a^2}| + C$$

66
$$\int x\sqrt{x^2 - a^2} dx = \frac{1}{3} (x^2 - a^2)^{\frac{3}{2}} + C$$

67
$$\int \frac{\sqrt{x^2 - a^2}}{x} dx = \sqrt{x^2 - a^2} + a \sin^{-1} \left(\frac{a}{x} \right) + C$$

68
$$\int \frac{\sqrt{x^2 - a^2}}{x^2} dx = -\frac{\sqrt{x^2 - a^2}}{x} + \ln|x + \sqrt{x^2 - a^2}| + C$$

69
$$\int \frac{dx}{\sqrt{x^2 - a^2}} = \ln|x + \sqrt{x^2 - a^2}| + C$$

70
$$\int \frac{xdx}{\sqrt{x^2 - a^2}} = \sqrt{x^2 - a^2} + C$$

71
$$\int \frac{x^2 dx}{\sqrt{x^2 - a^2}} = \frac{x}{2} \sqrt{x^2 - a^2} + \frac{a^2}{2} \ln|x + \sqrt{x^2 - a^2}| + C$$

72
$$\int \frac{dx}{x\sqrt{x^2 - a^2}} = -\frac{1}{a} \sin^{-1} \left(\frac{a}{x} \right) + C$$

73
$$\int \frac{dx}{(x+a)\sqrt{x^2 - a^2}} = \frac{1}{a} \sqrt{\frac{x-a}{x+a}} + C$$

74
$$\int \frac{dx}{(x-a)\sqrt{x^2 - a^2}} = -\frac{1}{a} \sqrt{\frac{x+a}{x-a}} + C$$

75
$$\int \frac{dx}{x^2\sqrt{x^2 - a^2}} = \left(\frac{\sqrt{x^2 - a^2}}{a^2 x} \right) + C$$

76
$$\int \frac{dx}{(x^2 - a^2)^{\frac{3}{2}}} = -\frac{x}{a^2\sqrt{x^2 - a^2}} + C$$

77
$$\int (x^2 - a^2)^{\frac{3}{2}} dx = -\frac{x}{8} (2x^2 - 5a^2) \sqrt{x^2 - a^2} + \frac{3a^4}{8} \ln|x + \sqrt{x^2 - a^2}| + C$$

78
$$\int \sqrt{a^2 - x^2} dx = \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \left(\frac{x}{a} \right) + C$$

79
$$\int x\sqrt{a^2 - x^2} dx = -\frac{1}{3} (a^2 - x^2)^{\frac{3}{2}} + C$$

80
$$\int x^2 \sqrt{a^2 - x^2} dx = \frac{x}{8} (2x^2 - a^2) \sqrt{a^2 - x^2} + \frac{a^4}{8} \sin^{-1} \left(\frac{x}{a} \right) + C$$

81
$$\int \frac{\sqrt{a^2 - x^2}}{x} dx = \sqrt{a^2 - x^2} + a \ln \left| \frac{x}{a + \sqrt{a^2 - x^2}} \right| + C$$

82
$$\int \frac{\sqrt{a^2 - x^2}}{x^2} dx = -\frac{\sqrt{a^2 - x^2}}{x} - \frac{\sin^{-1} x}{a} + C$$

83
$$\int \frac{dx}{\sqrt{1 - x^2}} = \sin^{-1} x + C$$

$$\int \frac{dx}{\sqrt{a^2 - x^2}} = \frac{\sin x}{a} + C$$

$$85 \quad \int \frac{x dx}{\sqrt{a^2 - x^2}} = -\sqrt{a^2 - x^2} + C$$

$$86 \quad \int \frac{x^2 dx}{\sqrt{a^2 - x^2}} = -\frac{x}{2}\sqrt{a^2 - x^2} + \frac{a^2}{2}\sin^{-1}\left(\frac{x}{a}\right) + C$$

$$87 \quad \int \frac{dx}{(x+a)\sqrt{a^2 - x^2}} = -\frac{1}{2}\sqrt{\frac{a-x}{a+x}} + C$$



[DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE](#)

$$88 \quad \int \frac{dx}{(x-a)\sqrt{a^2 - x^2}} = -\frac{1}{\sqrt{\frac{a+x}{a-x}}} + C$$

$$89 \quad \int \frac{dx}{(x+b)\sqrt{a^2 - x^2}} = \frac{1}{\sqrt{b^2 - a^2}} \frac{\sin^{-1}(bx + a^2)}{a(x+b)} + C, b > a.$$

$$90 \quad \int \frac{dx}{(x+b)\sqrt{a^2 - x^2}} = \frac{1}{\sqrt{a^2 - b^2}} \ln \left| \frac{x+b}{\sqrt{a^2 - b^2}\sqrt{a^2 - x^2} + a^2 + bx} \right| + C, b < a.$$

$$91 \quad \int \frac{dx}{x^2\sqrt{a^2 - x^2}} = -\frac{\sqrt{a^2 - x^2}}{a^2 x} + C$$

$$92 \quad \int (a^2 - x^2)^{\frac{3}{2}} dx = \frac{x}{8}(5a^2 - 2x^2)\sqrt{a^2 - x^2} + \frac{3a^4}{8}\sin^{-1}\left(\frac{x}{a}\right) + C$$

$$93 \quad \int \frac{dx}{(a^2 - x^2)^{\frac{3}{2}}} = \frac{x}{a^2\sqrt{a^2 - x^2}} + C$$

$$94 \quad \int \sin x dx = -\cos x + C$$

$$95 \quad \int \cos x dx = \sin x + C$$



[DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE](#)

$$96 \quad \int \sin^2 x dx = \frac{x}{2} - \frac{1}{4}\sin 2x + C$$

$$97 \quad \int \cos^2 x dx = \frac{x}{2} + \frac{1}{4}\sin 2x + C$$

$$98 \quad \int \sin^3 x dx = \frac{1}{3}\cos^3 x - \cos x + C = \frac{1}{12}\cos 3x - \frac{3}{4}\cos x + C$$

$$99 \quad \int \cos^3 x dx = \sin x - \frac{1}{3}\sin^3 x + C = \frac{1}{12}\sin 3x + \frac{3}{4}\sin x + C$$

$$100 \quad \int \frac{dx}{\cos x} = \int \sec x dx = \ln \left| \tan \left(\frac{x}{2} + \frac{\pi}{4} \right) \right| + C$$

$$101 \quad \int \frac{dx}{\sin x} = \int \csc x dx = \ln \left| \frac{\tan x}{2} \right| + C$$

$$102 \quad \int \frac{dx}{\sin^2 x} = \int \csc^2 x dx = -\cot x + C$$



$$103 \quad \int \frac{dx}{\cos^2 x} = \int \sec^2 x dx = \tan x + C$$



[DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE](#)

$$104 \quad \int \frac{dx}{\sin^3 x} = \int \csc^3 x dx = -\frac{\cos x}{2\sin^2 x} = \frac{1}{2}\ln \left| \frac{\tan x}{2} \right| + C$$

$$105 \quad \int \frac{dx}{\cos^3 x} = \int \sec^3 x dx = \frac{\sin x}{2\cos^2 x} + \frac{1}{2}\ln \left| \tan \left(\frac{x}{2} + \frac{\pi}{4} \right) \right| + C$$

106	$\int \sin x \cos x dx = -\frac{1}{4} \cos 2x + C$
107	$\int \sin^2 x \cos x dx = \frac{1}{3} \sin^3 x + C$
108	$\int \sin x \cos^2 x dx = -\frac{1}{3} \cos^3 x + C$
109	$\int \sin^2 x \cos^2 x dx = \frac{x}{8} - \frac{1}{32} \sin 4x + C$
110	$\int \tan x dx = -\ln \cos x + C$
111	$\int \frac{\sin x}{\cos^2 x} dx = \frac{1}{\cos x} + C = \sec x + C$
 पढ़ना हुआ आसान	DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE
112	$\int \frac{\sin^2 x}{\cos x} dx = \ln\left \tan\left(\frac{x}{2} + \frac{\pi}{4}\right)\right - \sin x + C$
113	$\int \tan^2 x dx = \tan x - x + C$
114	$\int \cot x dx = \ln \sin x + C$
115	$\int \frac{\cos x}{\sin^2 x} dx = -\frac{1}{\sin x} + C = -\operatorname{cosec} x + C$
116	$\int \frac{\cos^2 x}{\sin x} dx = \ln\left \frac{\tan x}{2}\right + \cos x + C$
117	$\int \cot^2 x dx = -\cot x - x + C$
118	$\int \frac{dx}{\cos x \sin x} = \ln \tan x + C$
119	$\int \frac{dx}{\sin^2 x \cos x} = -\frac{1}{\sin x} + \ln\left \tan\left(\frac{x}{2} + \frac{\pi}{4}\right)\right + C$
 पढ़ना हुआ आसान	DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE
120	$\int \frac{dx}{\sin x \cos^2 x} = \frac{1}{\cos x} + \ln\left \frac{\tan x}{2}\right + C$
121	$\int \frac{dx}{\sin^2 x \cos^2 x} = \tan x - \cot x + C$
122	$\int \sin mx \sin nx dx = -\frac{\sin(m+n)x}{2(m+n)} + \frac{\sin(m-n)x}{2(m-n)} + C, m^2 \neq n^2,$
123	$\int \sin mx \cos nx dx = -\frac{\cos(m+n)x}{2(m+n)} - \frac{\cos(m-n)x}{2(m-n)} + C, m^2 \neq n^2.$
124	$\int \cos mx \cos nx dx = \frac{\sin(m+n)x}{2(m+n)} + \frac{\sin(m-n)x}{2(m-n)} + C, m^2 \neq n^2.$
125	$\int \sec x \tan x dx = \sec x + C$
126	$\int \operatorname{cosec} x \cot x dx = -\operatorname{cosec} x + C$
127	$\int \sin x \cos^n x dx = -\frac{\cos^{n+1} x}{n+1} + C$
 पढ़ना हुआ आसान	DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE
128	

$$\int \sin x \cos^n x dx = -\frac{\sin^{n+1} x}{n+1} + C$$

$$129 \quad \int \sin^{-1} x dx = x \sin^{-1} x + \sqrt{1-x^2} + C$$

$$130 \quad \int \cos^{-1} x dx = x \cos^{-1} x - \sqrt{1-x^2} + C$$

$$131 \quad \int \tan^{-1} x dx = x \tan^{-1} x - \frac{1}{2} \ln(x^2 + 1) + C$$

$$132 \quad \int \cot^{-1} x dx = x \cot^{-1} x + \frac{1}{2} \ln(x^2 + 1) + C$$

$$133 \quad \int \sinh x dx = \cosh x + C$$

$$134 \quad \int \cosh x dx = \sinh x + C$$

$$135 \quad \int \tanh x dx = \ln \cosh x + C$$



[DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE](#)

$$136 \quad \int \coth x dx = \ln|\sinh x| + C$$

$$137 \quad \int \operatorname{sech}^2 x dx = \tanh x + C$$

$$138 \quad \int \operatorname{cosech}^2 x dx = -\coth x + C$$

$$139 \quad \int \operatorname{sech} x \tanh x dx = -\operatorname{sech} x + C$$

$$140 \quad \int \operatorname{cosech} x \coth x dx = -\operatorname{cosech} x + C$$

$$141 \quad \int e^x dx = e^x + C$$

$$142 \quad \int a^x dx = \frac{a^x}{\ln a} + C$$

$$143 \quad \int e^{ax} dx = \frac{e^{ax}}{a} + C$$



[DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE](#)

$$144 \quad \int x e^{ax} dx = \frac{e^{ax}}{a^2} (ax - 1) + C$$

$$145 \quad \int \ln x dx = x \ln x - x + C$$

$$146 \quad \int \frac{dx}{x \ln x} = \ln|\ln x| + C$$

$$147 \quad \int x^n \ln x dx = x^{n+1} \left[\frac{\ln x}{n+1} - \frac{1}{(n+1)^2} \right] + C$$

$$148 \quad \int e^{ax} \sin bx dx = \frac{a \sin bx - b \cos bx}{a^2 + b^2} e^{ax} + C$$

$$149 \quad \int e^{ax} \cos bx dx = \frac{a \cos bx + b \sin bx}{a^2 + b^2} e^{ax} + C$$

$$150 \quad \int x^m e^{mx} dx = \frac{1}{m} x^m e^{mx} - \frac{m}{m} \int x^{m-1} e^{mx} dx$$

$$151 \quad \int \frac{e^{mx}}{x^n} dx = -\frac{e^{mx}}{(n-1)x^{n-1}} + \frac{m}{n-1} \int \frac{e^{mx}}{x^{n-1}} dx, n \neq 1,$$

$$152 \quad \int \sinh^n x dx = \frac{1}{n} \sinh^{n-1} x \cosh x - \frac{n-1}{n} \int \sinh^{n-2} x dx$$

$$153 \quad \int \frac{dx}{\sinh^n x} = \frac{\cosh x}{(n-1)\sinh^{n-1} x} - \frac{n-2}{n-1} \int \frac{dx}{\sinh^{n-2} x}, n \neq 1.$$

$$154 \quad \int \cosh^n x dx = \frac{1}{n} \sinh x \cosh^{n-1} x \cosh x - \frac{n-1}{n} \int \cosh^{n-2} x dx$$

$$155 \quad \int \frac{dx}{\cosh^n x} = \frac{\sinh x}{n-1(\cosh^{n-1} x) + \frac{n-2}{n-1}} \int \frac{dx}{\cosh^{n-2} x}, n \neq 2.$$

$$156 \quad \int \sinh^n x \cosh^m x dx = \frac{\sinh x}{n+1 \frac{x \cosh^{m-1} x}{n+m} - \frac{m-1}{n+m} \int \sinh^n x \cosh^{m-2} x dx}$$

$$157 \quad \int \sinh nx \cosh^m x dx = \frac{\sinh^{n-1} x \cosh^{m+1} x}{n+m} - \frac{n-1}{n+m} \int \sinh^{n-2} x \cosh^m x dx$$

$$158 \quad \int \tanh^n x dx = -\frac{1}{n-1} \tanh^{n-1} x + \int \tanh^{n-2} x dx, n \neq 1.$$

$$159 \quad \int \coth^n x dx = -\frac{1}{n-1} \coth^{n-1} x + \int \coth^{n-2} x dx, n \neq 1.$$

$$160 \quad \int \operatorname{sech}^n x dx = \frac{\operatorname{sech}^{n-2} x \tanh x}{n-1} + \frac{n-2}{n-1} \int \operatorname{sech}^{n-2} x dx, n \neq 1.$$

$$161 \quad \int \sin^n x dx = -\frac{1}{n} \sin^{n-1} x \cos x + \frac{n-1}{n} \int \sin^{n-2} x dx$$

$$162 \quad \int \frac{dx}{\sin^n x} = -\frac{\cos x}{(n-1)\sin^{n-1} x} + \frac{n-2}{n-1} \int \frac{dx}{\sin^{n-2} x}, n \neq 1.$$

$$163 \quad \int \cos^n x dx = \frac{1}{n} \sin x \cos^{n-1} x + \frac{n-1}{n} \int \cos^{n-2} x dx$$

$$164 \quad \int \frac{dx}{\cos^n x} = \frac{\sin x}{(n-1)\cos^{n-1} x} + \frac{n-2}{n-1} \int \frac{dx}{\cos^{n-2} x}, n \neq 1$$

$$165 \quad \int \sin^n x \cos^m x dx = \frac{\sin^{n+1} x \cos^{m-1} x}{n+m} + \frac{m-1}{n+m} \int \sin^n x \cos^{m-2} x dx$$

$$166 \quad \int \sin^n x \cos^m x dx = -\frac{\sin^{n-1} x \cos^{m+1} x}{n+m} + \frac{n-1}{n+m} \int \sin^{n-2} x \cos^m x dx$$

$$167 \quad \int \tan^n x dx = \frac{1}{n-1} \tan^{n-1} x - \int \tan^{n-2} x dx, n \neq 1.$$

$$168 \quad \int \cot^n x dx = -\frac{1}{n-1} \cot^{n-1} x - \int \cot^{n-2} x dx, n \neq 1.$$


$$169 \quad \int \sec^n x dx = \frac{\sec^{n-2} x \tan x}{n-1} + \frac{n-2}{n-1} \int \sec^{n-2} x dx, n \neq 1.$$

$$170 \quad \int \operatorname{cosec}^n x dx = \frac{\operatorname{cosec}^{n-2} x \cot x}{n-1} + \frac{n-2}{n-1} \int \operatorname{cosec}^{n-2} x dx, n \neq 1$$

$$171 \quad \int x^n \ln^m x dx = \frac{x^{n+1} \ln^m x}{n+1} - \frac{m}{n+1} \int x^n \ln^{m-1} x dx$$

$$172 \quad \int \frac{\ln^m x}{x^n} dx = -\frac{\ln^m x}{(n-1)x^{n-1}} + \frac{m}{n-1} \int \frac{\ln^{m-1} x}{x^n} dx, n \neq 1.$$

$$173 \quad \int \ln^n x dx = x \ln^n x - n \int \ln^{n-1} x dx.$$

174	$\int x^n \sinh x dx = x^n \cosh x - n \int x^{n-1} \cosh x dx.$
175	$\int x^n \cosh x dx = x^n \sinh x - n \int x^{n-1} \sinh x dx.$
 पढ़ना हुआ आसान	DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE
176	$\int x^n \sin x dx = -x^n \cos x + n \int x^{n-1} \cos x dx$
177	$\int x^n \cos x dx = x^n \sin x - n \int x^{n-1} \sin x dx.$
178	$\int x^n \sin^{-1} x dx = \frac{x^{n+1}}{n+1} \sin^{-1} x - \frac{1}{n+1} \int \frac{x^{n+1}}{\sqrt{1-x^2}} dx$
179	$\int x^n \cos^{-1} x dx = \frac{x^{n+1}}{n+1} \cos^{-1} x + \frac{1}{n+1} \int \left(\frac{x^{n+1}}{\sqrt{1-x^2}} dx. \right.$
180	$\int x^n \tan^{-1} x dx = \frac{x^{n+1}}{n+1} \tan^{-1} x - \frac{1}{n+1} \int \frac{x^{n+1}}{1+x^2} dx.$
181	$\int \frac{x^n dx}{ax^n + b} = \frac{x}{a} - \frac{b}{a} \int \frac{dx}{ax^n + b}$
182	$\int \frac{dx}{(ax^2 + bx + c)^n} = \frac{-2ax - b}{(n-1)(b^2 - 4ac)(ax^2 + bx + c)^{n-1}} - \frac{2(2n-3)a}{(n-1)(b^2 - 4ac)} \int \frac{dx}{(ax^2 + bx + c)^{n-1}}, n \neq 1.$
183	$\int \frac{dx}{(x^2 + a^2)^n} = \frac{x}{2(n-1)a^2(x^2 + a^2)^{n-1}} + \frac{2n-3}{2(n-1)a^2} \int \frac{dx}{(x^2 + a^2)^{n-1}}, n \neq 1.$
 पढ़ना हुआ आसान	DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE
184	$\int \frac{dx}{(x^2 - a^2)^n} = -\frac{x}{2(n-1)a^2(x^2 - a^2)^{n-1}} - \frac{2n-3}{2(n-1)a^2} \int \frac{dx}{(x^2 - a^2)^{n-1}}, n \neq 1.$
185	$\int_a^b f(x) dx = \lim_{x \rightarrow \infty} \sum_{i=1}^n f(\xi_i) \Delta x_i, \text{ where } \Delta x_i = x_i - x_{i-1}, x_{i-1} \leq \xi_i \leq x_i.$
186	$\int_a^b 1 dx = b - a$
187	$\int_a^b kf(x) dx = k \int_a^b f(x) dx$
188	$\int_a^b [f(x) + g(x)] dx = \int_a^b f(x) dx + \int_a^b g(x) dx$
189	$\int_a^b [f(x) - g(x)] dx = \int_a^b f(x) dx - \int_a^b g(x) dx$
190	$\int_a^a f(x) dx = 0$
191	$\int_a^b f(x) dx = - \int_b^a f(x) dx$
 पढ़ना हुआ आसान	DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE
192	$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx \text{ if } a < c < b.$
193	$\int_a^b f(x) dx \geq 0 \text{ if } f(x) \geq 0 \text{ on } [a, b]$
194	$\int_a^b f(x) dx \leq 0 \text{ if } f(x) \leq 0 \text{ on } [a, b]$
195	Fundamental Theorem of Calculus

$$\int_a^b f(x)dx = F(x) \Big|_a^b = F(b) - F(a) \text{ if } F'(x) = f(x).$$

Method of substitution

196 **If** $x = g(t)$, **then** $\int_a^b f(x)dx = \int_c^d f(g(t))g'(t)dt$, **where** $c = g^{-1}(a)$, $d = g^{-1}(b)$.

Integration by Parts

197 $\int_a^b u dv(uv) \Big|_a^b - \int_a^b v du$

Traezoidal Rule

198 $\int_a^b f(x)dx = \frac{b-a}{2n} \left[f(x_0) + f(x_n) + 2 \sum_{i=1}^{n-1} f(x_i) \right]$

Simpson's Rule

199 $\int_a^b f(x)dx = \frac{b-a}{3n} [f(x_0) + 4f(x_1) + 2f(x_2) + 4f(x_3) + \dots + 2f(x_4) + \dots + 4f(x_{n-1}) + f(x_n)]$, **where** $x_i = a + \frac{b-a}{n}i$, $i, 0, 1, 2, \dots, n$.



[DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE](#)

Area Under a Curve

200 $S = \int_a^b f(x)dx = F(b) - F(a)$, **where** $F'(x) = f(x)$.

Area Between Two Curves

201 $S = \int_a^b [f(x) - g(x)]dx = F(b) - G(b) - F(a) + G(a)$, **where** $F'(x) = f(x)$, $G'(x) = g(x)$.

202 **The definite integral** $\int_a^b f(x)dx$ **is called an improper integral if**, a **or** b **is infinite, $f(x)$ has one or more points of discontinuity in the interval** $[a, b]$.

203 **If** $f(x)$ **is a continuous functn on** $[a, \infty)$ **then** $\int_a^\infty f(x)dx = \lim_{n \rightarrow \infty} \int_a^n f(x)dx$.

204 **If** $f(x)$ **is continuous function on** $(-\infty, b)$, **then** $\int_{-\infty}^b f(x)dx = \lim_{n \rightarrow \infty} \int_n^b f(x)dx$.

205 $\int_{-\infty}^\infty f(x)dx = \int_{-\infty}^c f(x)dx + \int_c^\infty f(x)dx$

Comparison Theorem:

206 **Let** $f(x)$ **and** $g(x)$ **be continuous functions on the closed interval** $[a, \infty)$. **Suppose that** $0 \leq g(x) \leq f(x)$ **for all** x **in** $[a, \infty)$., **If** $\int_a^\infty f(x)dx$ **is convergent, then** $\int_a^\infty g(x)dx$ **is also convergent, if** $\int_a^\infty g(x)dx$ **is divergent then** $\int_a^\infty f(x)dx$ **is also divergent.**

Absolute Convergence

207 **If** $\int_a^\infty |f(x)|dx$ **is convergent then the integral** $\int_a^\infty f(x)dx$ **is absolutely convergent.**



[DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE](#)

Discontinuous Integrand

208 **Let** $f(x)$ **be a function which is continuous on the interval** $[a, b)$ **but is discontinuous at** $x = b$. **Then** $\int_a^b f(x)dx = \lim_{\xi \rightarrow 0^+} \int_a^{b-\xi} f(x)dx$.

209 **Let** $f(x)$ **be a continuous function for all real numbers** x **in the interval** $[a, b]$ **except for some point** c **in** (a, b) . **Then** $\int_a^b f(x)dx = \lim_{\xi \rightarrow 0^+} \int_a^{c-\xi} f(x)dx + \lim_{\delta \rightarrow 0^+} \int_{c+\delta}^b f(x)dx$

Double Integral Functions of two variables

$f(x, y), (u, v)$

Double integrals: $\iint_r f(x, y)dx dy, \iint_r g(x, y)dx dy$

Riemann sum: $\sum_{i=1}^m \sum_{j=1}^n (f(u_i, v_j) \Delta x_i \Delta y_j)$

Small changes: $\Delta x_i, \Delta y_j$

Regions of integration :R,S

Polar coordinates $r\theta$

Area:A, Surface area;

Volume of a solid:V

Mass of a lamina:m

Density; $\rho(x, y)$

First moments : M_x, M_y

Moments of inertial: I_x, I_y, I_0

Charge of a plate: Q

Charge density: $\sigma(x, y)$,

Coordinates of center of mass: \bar{x}, \bar{y}

Average of a function : μ

211
$$\iint_R [f(x, y) + g(x, y)]dA = \iint_R f(x, y)dA + \iint_R g(x, y)dA$$

212
$$\iint_R [f(x, y) - g(x, y)]dA = \iint_R f(x, y)dA - \iint_R g(x, y)dA$$

213
$$\iint_R kf(x, y)dA = k \iint_R (x, y)dA \text{ where } k \text{ is a constant.}$$

214 **If $f(x, y) \leq g(x, y)$ on \mathbb{R} , then $\iint_R f(x, y)dA \leq \iint_R g(x, y)dA$.**

215 **If $f(x, y) \geq 0$ on R and $S \subset \text{sec } R$, then $\iint_S f(x, y)dA \leq \iint_R f(x, y)dA$.**



[DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE](#)

216 **If $f(x, y) \geq 0$ on R and R and S are non-overlapping regions, then $\iint_{R \cup S} f(x, y)dA = \iint_R f(x, y)dA + \iint_S f(x, y)dA$ Here $R \cup S$ is the union of the regions R and S .**

217 **Integrated Integrals and Fubini's Theorem**

$$\iint_R f(x, y)dA = \int_a^b \int_{p(x)}^{q(x)} f(x, y)dydx \text{ for a region on type I, } R = \{(x, y) \mid a \leq x \leq b, p(x) \leq y \leq q(x)\}$$

218 **Doubles Integrals over Rectangular Regions**

If R is the rectangular region $[a, b] \times [c, d]$, then $\iint_R f(x, y)dxdy = \int_a^b \left(\int_c^d f(x, y)dy \right) dx = \int_c^d \left(\int_a^b f(x, y)dx \right) dy$. In the special case where the integrand $f(x, y)$ can be written as $g(x)h(y)$ we have $\iint_R f(x, y)dxdy = \iint_R g(x)h(y)dxdy = \left(\int_a^b g(x)dx \right) \left(\int_c^d h(y)dy \right)$

219 **Change of Variables**

$$\iint_R f(x, y)dxdy \equiv \iint_S f[x(u, v), y(u, v)] \left| \frac{\partial(x, y)}{\partial(u, v)} \right| dudv, \text{ where } \left| \frac{\partial(x, y)}{\partial(u, v)} \right| = \left| \begin{pmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{pmatrix} \right| \neq 0$$
 is the jacobian of the transformators
 $(x, y) \rightarrow (u, v)$ and S is the pullback of R which can be computed by $x = x(u, v), y = y(u, v)$ into the definitins of R .

220 **Polar Coordinates**

$x = r \cos \theta, y = r \sin \theta$

221 **Double Integrals in Polar Coordinates**

The differential $dxdy$ for Polar Coordinate is $dxdy = \left| \frac{\partial(x, y)}{\partial(r, \theta)} \right| drd\theta = r drd\theta$. Let the region R is determined as follows:
 $0 \leq g(\theta), r \leq h(\theta), \alpha \leq \theta \leq \beta$, where $\beta - \alpha \leq 2\pi$ Then $\iint_R f(x, y)dxdy = \int_\alpha^\beta \int_{g(\theta)}^{h(\theta)} f(r \cos \theta, r \sin \theta) r drd\theta$.

222 **Area of Region**

$$A = \int_a^b \int_{g(x)}^{f(x)} dydx \text{ (for a type I region).}$$

223 **Volume of a Solid**

$$V = \iint_R f(x, y) dA$$

Area and Volume in Polar Coordinates

224 **If S is a region in the xy-plane bounded by $\theta = \alpha, \theta = \beta, r = h(\theta), r = g(\theta)$, then** $A = \iint_S dA = \int_{\alpha}^{\beta} \int_{h(\theta)}^{g(\theta)} r dr d\theta, V = \iint_S f(r, \theta) r dr d\theta.$

225 **Surface Area** $S = \iint_R \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2} dx dy$

226 **Mass of a Lamina** $m = \iint_R \rho(x, y) dA,$
where the lamina occupies a regions R and its density at a point (x, y) is $\rho(x, y)$

Moments

227 **The moment of the lamna about the x-axis is given by formula** $M_x = \iint_R y \rho(x, y) dA.$ **The moment of the lamina about the y-axis is** $M_y = \iint_R x \rho(x, y) dA.$ **The moment of inertia about the x-axis is** $I_x = \iint_R y^2 \rho(x, y) dA.$ **The moment of inertia about the y-axis is** $I_y = \iint_R x^2 \rho(x, y) dA.$ **The polar moment of inertia is** $I_0 = \iint_R (x^2 + y^2) \rho(x, y) dA$

Center of Mass

228 $\bar{x} = \frac{M_y}{m} = \frac{1}{m} \iint_R x \rho(x, y) dA = \frac{\iint_R x \rho(x, y) dA}{\iint_R \rho(x, y) dA},$
 $\bar{y} = \frac{M_x}{m} = \frac{1}{m} \iint_R y \rho(x, y) dA = \frac{\iint_R y \rho(x, y) dA}{\iint_R \rho(x, y) dA},$

Charge of a Plate

229 $Q = \iint_R \sigma(x, y) dA,$ where electrical charge is distributed over a region R and its charge density at a point (x, y) is $\sigma(x, y)$

Average of Function

230 $\mu = \frac{1}{S} \iint_R f(x, y) dA,$ where $S = \iint_R dA.$

Definition of Triple Integral

231 **The triple integral over as parallelepiped $[a, b] \times [c, d] \times [r, s]$ is defined to be**
$$\iiint_{[a,b] \times [c,d] \times [r,s]} f(x, y, z) dV = \lim_{\substack{\max \Delta x_i \rightarrow 0 \\ \max \Delta y_j \rightarrow 0 \\ \max \Delta z_k \rightarrow 0}} \sum_{i=1}^m \sum_{j=1}^n \sum_{k=1}^p f(u_i, v_j, w_k) \Delta x_i \Delta y_j \Delta z_k, \text{ where } (u_i, v_j, w_k) \text{ is some point in the paralleloiped}$$

 $(x_{i-1}, x_i) \times (y_{j-1}, y_j) \times (z_{k-1}, z_k) \text{ and } \Delta x_i = x_i - x_{i-1}, \Delta y_j = y_j - y_{j-1}, \Delta z_k = z_k - z_{k-1}.$

232 $\iiint_G [f(x, y, z) + g(x, y, z)] dV = \iiint_G f(x, y, z) dV + \iiint_G g(x, y, z) dV$

233 $\iiint_G [f(x, y, z) - g(x, y, z)] dV = \iiint_G f(x, y, z) dV - \iiint_G g(x, y, z) dV$

234 $\iiint_G k f(x, y, z) dV = k \iiint_G f(x, y, z) dV$ where k is a constant.

235 **If $f(x, y, z) \geq 0$ and G and T are nonoverlapping basic regions then** $\iiint_{G \cup T} f(x, y, z) dV = \iiint_G f(x, y, z) dV + \iiint_T f(x, y, z) dV.$ **Here $G \cup T$ is the union of the regions G and T.**

Evaluation of Triple Integrals by Repeated Integrals

236 **If the solid G is the set of points (x, y, z) such that $(x, y) \in R, \chi_1(x, y) \leq z \leq \chi_2(x, y)$ then** $\iiint_G f(x, y, z) dx dy dz = \iint_R \left[\int_{\chi_1(x, y)}^{\chi_2(x, y)} f(x, y, z) dz \right] dx dy,$ **where R is projection of G onto the xy-plane. If the solid G is the set of points (x, y, z) such that $a \leq x \leq b, \varphi_1(x) \leq y \leq \varphi_2(x), \chi_1(x, y) \leq z \leq \chi_2(x, y),$ then**
$$\iiint_G f(x, y, z) dx dy dz = \int_a^b \left[\int_{\varphi_1(x)}^{\varphi_2(x)} \left(\int_{\chi_1(x, y)}^{\chi_2(x, y)} f(x, y, z) dz \right) dy \right] dx$$

Triple Integrals over Parallelepiped

237

If G is a parallelepiped $[a, b] \times [c, d] \times [r, s]$ then $\iiint_G f(x, y, z) dx dy dz = \int_a^b \left[\int_c^d \left(\int_r^s dz \right) dy \right] dx$. In the special case where in the integrand $f(x, y, z)$ can be written as $g(x)h(y)k(z)$ we have $\iiint_G f(x, y, z) dx dy dz = \left(\int_a^b g(x) dx \right) \left(\int_c^d h(y) dy \right) \left(\int_r^s k(z) dz \right)$.

Change of Variables

238 $\iiint_G f(x, y, z) dx dy dz = \iiint_S f(u, v, w, y(u, v, w), z(u, v, w)) \left| \frac{\partial(x, y, z)}{\partial(u, v, w)} \right| dx dy dz$, where $\left| \frac{\partial(x, y, z)}{\partial(u, v, w)} \right| = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} & \frac{\partial x}{\partial w} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} & \frac{\partial y}{\partial w} \\ \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} & \frac{\partial z}{\partial w} \end{vmatrix} \neq 0$ is the jacobian of the transformation $(x, y, z) \rightarrow (u, v, w)$ and S is the pull back of G which can be computed by $x = x(u, v, w), y = y(u, v, w), z = z(u, v, w)$ into the definition of G .

Triple Integrals in Spherical Coordinates

239 The Differential $dx dy dz$ for Spherical Coordinates is $dx dy dz = \left| \frac{\partial(x, y, z)}{\partial(r, \theta, \varphi)} \right| dr d\theta d\varphi = r^2 \sin \theta dr d\theta d\varphi$. $\iiint_R f(x, y, z) dx dy dz = \iiint_S f(r \sin \theta \cos \varphi, r \sin \theta \sin \varphi, r \cos \theta) r^2 \sin \theta dr d\theta d\varphi$, where the solid S is the pullback of G is spherical coordinates. the angle θ ranges from 0 to 2π , the angle φ ranges from $0 \rightarrow \pi$.



[DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE](#)

Volume of a solid

240 $V = \iiint_G dx dy dz$

Volume of Cylindrical Coordinates

241 $V = \iiint_{S(r, \theta, z)} r dr d\theta dz$

242 **Volume of Spherical Coordinates** $V = \iiint_{S(r, \theta, \varphi)} r^2 \sin \theta dr d\theta d\varphi$

243 **Mass of a solid** $m = \iiint_G \mu(x, y, z) dV$, where the solid occupies a region G and its density at a point (x, y, z) is $\mu(x, y, z)$

244 **Center of mass of a solid** $\bar{x} = \frac{M_{yz}}{m}, \bar{y} = \frac{M_{xz}}{m}, \bar{z} = \frac{M_{xy}}{m}$, where $M_{yz} = \iiint_G x \mu(x, y, z) dV, M_{xz} = \iiint_G y \mu(x, y, z) dV, M_{xy} = \iiint_G z \mu(x, y, z) dV$ are the first moments about the coordinate planes $x = 0, y = 0, z = 0$ respectively, $m(x, y, z)$ is the density function.

Moments of Inertia about the xy-plane (or $z = 0$), yz-plane ($x = 0$) and xz-plane ($y = 0$)

245 $I_{xy} = \iiint_G z^2 \mu(x, y, z) dV, I_{yz} = \iiint_G x^2 \mu(x, y, z) dV, I_{xz} = \iiint_G y^2 \mu(x, y, z) dV$.

Moment of Inertia about the x-axis, y-axis, and z-axis

246 $I_x = I_{xy} + \left(I_{xz} = \iiint_G (z^2 + y^2) \mu(x, y, z) dV \right)$
 $I_y = I_{xy} + \left(I_{yz} = \iiint_G (z^2 + x^2) \mu(x, y, z) dV, I_z = I_{xz} + \left(I_{yz} = \iiint_G (y^2 + x^2) \mu(x, y, z) dV \right) \right)$

Polar Moment of Inertia

247 $I_0 = I_{xy} + I_{yz} + I_{xz} = \iiint_G (x^2 + y^2 + z^2) \mu(x, y, z) dV$



[DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE](#)

Line Integral of a Scalar Function

248 Let a curve C be given by the vector function $\vec{r} = \vec{r}(s), 0 \leq s < +S$, and a scalar function F is defined over the curve C . Then $\int_0^S F(\vec{r}(s)) ds = \int_C F(x, y, z) ds = \int_C F ds$, where ds is the arc length differential.

249 $\int_{C_1 \cup C_2} F ds = \int_{C_1} F ds + \int_{C_2} F ds$

250 If the smooth Curve C is parametrized by $\vec{r} = \vec{r}(t), a \leq t \leq \beta$, then $\int_C F(x, y, z) ds = \int_a^\beta F(x(t), y(t), z(t)) \sqrt{(x'(t))^2 + (y'(t))^2 + (z'(t))^2} dt$.

251 If C is a smooth curve in the xy -plane given by the equation $y = f(x), a \leq x \leq b$, then $\int_C (x, y) ds = \int_a^b F(x, f(x)) \sqrt{1 + (f'(x))^2} dx$.

Line Integral of Scalar Function in Polar coordinates

252

$$\int_C F(x, y) ds = \int_\alpha^\beta F(r \cos \theta, r \sin \theta) \sqrt{r^2 - \left(\frac{dr}{d\theta}\right)^2} d\theta, \text{ where the curve C is defined by the polar function } r(\theta).$$

Line Integral of Vector Field

253

Let a curve C be defined by the vector function $\vec{r} = \vec{r}(s), 0 \leq s \leq S$. Then $\frac{d\vec{r}}{ds} = \vec{t} = (\cos \alpha, \cos \beta, \cos \gamma)$ is the unit vector of the tangent line to this curve.

Properties of Line Integrals of Vector fields

254

$\int_{-C} (\vec{F} \cdot d\vec{r}) = - \int_C (\vec{F} \cdot d\vec{r})$, where -C denote the curve with the opposite orientation.
 $\int_C (\vec{F} \cdot d\vec{r}) = \int_{C_1 \cup C_2} (\vec{F} \cdot d\vec{r}) = \int_{C_1} (\vec{F} \cdot d\vec{r}) + \int_{C_2} (\vec{F} \cdot d\vec{r})$, where C is the union of the curve C_1 and C_2 .

255

If the curve C is parameterized by $\vec{r}(t) = \{x(t), y(t), z(t)\}, \alpha \leq t \leq \beta$, then $\int_C P dx + Q dy + R dz = \int_\alpha^\beta \left(P(x(t), y(t), z(t)) \frac{dx}{dt} + Q(x(t), y(t), z(t)) \frac{dy}{dt} + R(x(t), y(t), z(t)) \frac{dz}{dt} \right) dt$.



[DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE](#)

256

If C lies in the xy-plane and given by the equation $y = f(x)$, then $\int_C P dx + Q dy = \int_a^b \left(P(x, f(x)) + Q(x, f(x)) \frac{df}{dx} \right) dx$.

Green's Theorem

257

$\iint_R \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx dy = \oint_C P dx + Q dy$, where $\vec{F} = P(x, y) \vec{i} + Q(x, y) \vec{j}$ is a continuous vector function with continuous first partial derivatives $\frac{\partial P}{\partial y}, \frac{\partial Q}{\partial x}$ in a some domain R, which is bounded by a close, piecewise smooth curve C.

258

Area of a Region R bounded by the curve C $S = \iint_R dx dy = \frac{1}{2} \oint_C x dy - y dx$

Path Independence of Line Integrals

259

The line integral of a vector function $\vec{F} = P \vec{i} + Q \vec{j} + R \vec{k}$ is said to be path independent, if and only if P, Q and R are continuous in a domain D, and if there exists some scalar function $u = u(x, y, z)$ (a scalar potential) in D such that $\vec{F} = \nabla u$, or $\frac{\partial u}{\partial x} = P, \frac{\partial u}{\partial y} = Q, \frac{\partial u}{\partial z} = R$. Then $\int_C \vec{F}(\vec{r}) \cdot d\vec{r} = \int_C P dx + Q dy + R dz = u(B) - u(A)$.

260

Test for a conservative field

A vector field of the form $\vec{F} = \nabla u$ is called a conservative field. The line integral of a vector function $\vec{F} = P \vec{i} + Q \vec{j} + R \vec{k}$ is path independent if and only if $\text{curl } \vec{F} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ P & Q & R \end{vmatrix} = \vec{0}$. If the line integral is taken in xy-plane so that $\int_C P dx + Q dy = (B) - u(A)$, then the test for determining if a vector field is conservative can be written in the form $\frac{\partial P}{\partial y} = \frac{\partial Q}{\partial x}$.

Length of a Curve

261

$L = \int_C ds = \int_\alpha^\beta \left| \frac{d\vec{r}}{dt}(t) \right| dt = \int_\alpha^\beta \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 + \left(\frac{dz}{dt}\right)^2} dt$, where C is a piecewise smooth curve described by the position vector $\vec{r}(t), \alpha \leq t \leq \beta$. If the curve C is two dimensional, then $L = \int_C ds = \int_\alpha^\beta \left| \frac{d\vec{r}}{dt}(t) \right| dt = \int_\alpha^\beta \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$. The curve C is the graph of a function $y = f(x)$ in the xy-plane ($a \leq x \leq b$), then $L = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$

Length of a curve in polar coordinates


262



$L = \int_\alpha^\beta \sqrt{\left(\frac{dr}{d\theta}\right)^2 + r^2} d\theta$, where the curve C is given by the equation $r = r(\theta), \alpha \leq \theta \leq \beta$ in polar coordinates.

Mass of a wire

263

$m = \int_C \rho(x, y, z) ds$, where $\rho(x, y, z)$ is the mass per unit length of the wire. If C is a curve parameterized by the vector function $\vec{r}(t) = \{x(t), y(t), z(t)\}$, then the mass can be computed by the formula $m = \int_\alpha^\beta \rho \left(x(t), y(t), z(t) \right) \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 + \left(\frac{dz}{dt}\right)^2} dt$. If C is a curve in xy-plane then the mass of the wire is given by $m = \int_C \rho(x, y, z) ds$, or $m = \int_\alpha^\beta \rho \left(x(t), y(t) \right) \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$ (in parametric form)

264	<p>Center of mass of a wire</p> $\bar{x} = \frac{M_{yz}}{m}, \bar{y} = \frac{M_{xz}}{m}, \bar{z} = \frac{M_{xy}}{m} \text{ where } M_{yz} = \int_C x\rho(x, y, z) ds, M_{xz} = \int_C y\rho(x, y, z) ds, M_{xy} = \int_C z\rho(x, y, z) ds,$
265	<p>Moments of Inertia</p> <p>The moments of inertia about the x-axis, y-axis, and z-axis are given by the formulas</p> $I_x = \int_C (y^2 + z^2)\rho(x, y, z) ds, I_y = \int_C (x^2 + z^2)\rho(x, y, z) ds, I_z = \int_C (x^2 + y^2)\rho(x, y, z) ds$
266	<p>Area of a Region Bounded by a closed curve</p> $S = \oint_C x dy - \int_C y dx = \frac{1}{2} \oint_C x dy - y dx.$
267	<p>Volume of a Solid Formed by Rotating a Closed Curve about the x-axis</p> $V = -\pi \int_C y^2 dx = -2\pi \int_C xy dy = -\frac{\pi}{2} \int_C 2xy dy + y^2 dx$
268	<p>Work done by a force</p> <p>\vec{F} on an object moving along a curve C is given by the line integral $W = \int_C \vec{F} \cdot d\vec{r}$, where \vec{F} is the vector force field acting on the object $d\vec{r}$ is the unit tangent vector.</p>
269	<p>Ampere's Law</p> $\oint_C \vec{B} \cdot d\vec{r} = \mu_o I.$ <p>The line integral of a magnetic field \vec{B} around a closed path C is equal to the total current I flowing through the area bounded by the path.</p>
270	<p>Faraday's Law</p> $\epsilon = \int_C \vec{E} \cdot d\vec{r} = -\frac{d\psi}{dt}$ <p>The electromotive force (emf) epsilon induced around a closed loop C is equal to the rate to the change of magnetic flux ψ passing through the loop.</p>
271	<p>Surface Integral of a scalar function</p> <p>Let a surface S be given by the position vector $\vec{r}(u, v) = x(u, v)\vec{i} + y(u, v)\vec{j} + z(u, v)\vec{k}$, where (u, v) ranges over some domain $D(u, v)$ of the uv-plane.</p> <p>The surface integral of a scalar function $f(x, y, z)$ over the surface S is defined as $\iint_S f(x, y, z) dS = \iint_{D(u, v)} f(x(u, v), y(u, v), z(u, v)) \left \frac{\partial \vec{r}}{\partial u} \times \frac{\partial \vec{r}}{\partial v} \right du dv$,</p> <p>where the partial derivatives $\frac{\partial \vec{r}}{\partial u}$ and $\frac{\partial \vec{r}}{\partial v}$ are given by</p> $\frac{\partial \vec{r}}{\partial u} = \frac{\partial x}{\partial u}(u, v)\vec{i} + \frac{\partial y}{\partial u}(u, v)\vec{j} + \frac{\partial z}{\partial u}(u, v)\vec{k}, \frac{\partial \vec{r}}{\partial v} = \frac{\partial x}{\partial v}(u, v)\vec{i} + \frac{\partial y}{\partial v}(u, v)\vec{j} + \frac{\partial z}{\partial v}(u, v)\vec{k} \text{ and } \frac{\partial \vec{r}}{\partial u} \times \frac{\partial \vec{r}}{\partial v} \text{ is the cross product.}$
 DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE	
272	<p>If the surface S is given by the equation $z = z(x, y)$ where $z(x, y)$ is a differentiable function in the domain $D(x, y)$, then</p> $\iint_S f(x, y, z) dS = \iint_{D(x, y)} f(x, y, z(x, y)) \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2} dx dy.$
273	<p>Surface integral of the vector field \vec{F} over the surface S If S is oriented outward, then</p> $\iint_S \vec{F}(x, y, z) \cdot d\vec{S} = \iint_S \vec{F}(x, y, z) \cdot \vec{n} dS = \iint_{D(u, v)} \vec{F}(x(u, v), y(u, v), z(u, v)) \cdot \left[\frac{\partial \vec{r}}{\partial u} \times \frac{\partial \vec{r}}{\partial v} \right] du dv.$
274	<p>If S is oriented inward, then $\iint_S \vec{F}(x, y, z) \cdot d\vec{S} = \iint_S \vec{F}(x, y, z) \cdot \vec{n} dS = \iint_D (u, v) \vec{F}(x(u, v), y(u, v), z(u, v)) \cdot \left[\frac{\partial \vec{r}}{\partial u} \times \frac{\partial \vec{r}}{\partial v} \right] du dv. d\vec{S} = \vec{n} dS$</p> <p>is called the vector element of the surface. Dot means the scalar product of the appropriate vectors. The partial derivatives $\frac{\partial \vec{r}}{\partial u}$ and $\frac{\partial \vec{r}}{\partial v}$ are given by</p> $\frac{\partial \vec{r}}{\partial u} = \frac{\partial x}{\partial u}(u, v)\vec{i} + \frac{\partial y}{\partial u}(u, v)\vec{j} + \frac{\partial z}{\partial u}(u, v)\vec{k}, \frac{\partial \vec{r}}{\partial v} = \frac{\partial x}{\partial v}(u, v)\vec{i} + \frac{\partial y}{\partial v}(u, v)\vec{j} + \frac{\partial z}{\partial v}(u, v)\vec{k}.$
275	<p>If the surface S is given by the equation $z = z(x, y)$ where $z(x, y)$ is a differentiable function in the domain $D(x, y)$, then If S is oriented upward, i.e. the k-th component of the normal vector is positive, then $\int_S \text{vec} F(x, y, z) \cdot d\text{vec} S = \int_{D(x, y)} \text{vec} F(x, y, z) \cdot \left(\frac{\partial z}{\partial x} \vec{i} + \frac{\partial z}{\partial y} \vec{j} + \vec{k} \right) dx dy$. If S is oriented downward, i.e. the k-th component of the normal vector is negative, then $\int_S \text{vec} F(x, y, z) \cdot d\text{vec} S = \int_{D(x, y)} \text{vec} F(x, y, z) \cdot \left(\frac{\partial z}{\partial x} \vec{i} + \frac{\partial z}{\partial y} \vec{j} - \vec{k} \right) dx dy$.</p>
276	$\iint_S (\vec{F} \cdot \vec{n}) dS = \iint_S P dy dz + Q dz dx + R dx dy = \iint_S (P \cos \alpha + Q \cos \beta + R \cos \gamma) dS,$ <p>where $P(x, y, z), Q(x, y, z), R(x, y, z)$ are the components of the vector field \vec{F}, $\cos \alpha, \cos \beta, \cos \gamma$ are the angles between the outer unit normal vector \vec{n} and the x-axis, y-axis, and z-axis, respectively.</p>
277	

	<p>If The surface S is given in parametric form by the vector $\vec{r}(x(u, v), y(u, v), z(u, v))$, then the latter formula can be written as</p> $\iint_S (\vec{F} \cdot \vec{n}) dS = \iint_S P dydz + Q dzdx + R dx dy = \iint_{D(u, v)} \left (P, Q, R), \begin{matrix} \frac{\partial x}{\partial u}, \frac{\partial y}{\partial u}, \frac{\partial z}{\partial u} \\ \frac{\partial x}{\partial v}, \frac{\partial y}{\partial v}, \frac{\partial z}{\partial v} \end{matrix} \right dudv, \text{ where } (u, v) \text{ ranges over some domain } D(u, v) \text{ of the } uv\text{-plane.}$
278	<p>Divergence Theorem</p> $\oiint_S \vec{F} \cdot d\vec{S} = \iiint_G (\nabla \cdot \vec{F}) dV, \text{ where } \vec{F}(x, y, z) = \{P(x, y, z), Q(x, y, z), R(x, y, z)\} \text{ is a vector field whose components } P, Q, \text{ and } R \text{ have continuous partial derivatives, } \nabla \cdot \vec{f} = \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z} \text{ is the divergence of } \vec{F} \text{ also denoted } \operatorname{div} \vec{F}. \text{ The symbol } \oiint \text{ indicates that the surface integral is taken over a closed surface.}$
279	<p>Divergence Theorem in Coordinate Form</p> $\oiint_S P dydz + Q dzdx + R dx dy = \iiint_G \left(\frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z} \right) dx dy dz.$
	 DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE
280	<p>Stoke's Theorem</p> $\oint_C \vec{F} \cdot d\vec{r} = \iint_S (\nabla \times \vec{F}) \cdot d\vec{S}, \text{ where } \vec{F}(x, y, z) = \{P(x, y, z), Q(x, y, z), R(x, y, z)\} \text{ is a vector field whose components } P, Q, \text{ and } R \text{ have continuous partial derivatives } \nabla \times \vec{f} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ P & Q & R \end{vmatrix} = \left(\frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z} \right) \vec{i} + \left(\frac{\partial P}{\partial z} - \frac{\partial R}{\partial x} \right) \vec{j} + \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) \vec{k} \text{ is the curl of } \vec{F}, \text{ also denoted } \operatorname{curl} \vec{F}. \text{ The symbol, } \oint \text{ indicates that the line integral is taken over a closed curve.}$
281	<p>Stroke's Theorem in Coordinate Form</p> $\oint_C P dx + Q dy + R dz = \iint_S \left(\frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z} \right) dy dz + \left(\frac{\partial P}{\partial z} - \frac{\partial R}{\partial x} \right) dz dx + \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx dy$
282	<p>Surface Area $A = \iint_S dS$</p>
283	<p>If the surface S is parametrized by the vector $\vec{r}(u, v) = x(u, v)\vec{i} + y(u, v)\vec{j} + z(u, v)\vec{k}$, then the surface area is $A = \iint_{D(u, v)} \left \frac{\partial \vec{r}}{\partial u} \times \frac{\partial \vec{r}}{\partial v} \right dudv, \text{ where } D(u, v) \text{ is the domain where the surface } \vec{r}(u, v) \text{ is defined.}$</p>
284	<p>If S given explicitly by the function $z(x, y)$ then the surface area is $A = \iint_{D(x, y)} \sqrt{1 + \left(\frac{\partial z}{\partial x} \right)^2 + \left(\frac{\partial z}{\partial y} \right)^2} dx dy, \text{ where } D(x, y) \text{ is the projection of the surface } S \text{ onto the } xy\text{-plane.}$</p>
285	<p>Mass of a surface $m = \iint_S \mu(x, y, z) dS, \text{ where } \mu(x, y, z) \text{ is the mass per unit area (density function).}$</p>
286	<p>Center of Mass of a shell $\bar{x} = \frac{M_{yz}}{m}, \bar{y} = \frac{M_{xz}}{m}, \bar{z} = \frac{M_{xy}}{m}, \text{ where } M_{yz} = \iint_S x\mu(x, y, z) dS, M_{xz} = \iint_S y\mu(x, y, z) dS, M_{xy} = \iint_S z\mu(x, y, z) dS, \text{ are the first moments about the coordinate planes } x=0, y=0, z=0, \text{ respectively. } \mu(x, y, z) \text{ is the density function.}$</p>
287	<p>Moment of inertia the xy-plane (or z=0), yz-plane (x=0), and xz-plane (y=0) $I_{xy} = \iint_S z^2 \mu(x, y, z) dS, I_{yz} = \iint_S x^2 \mu(x, y, z) dS, I_{xz} = \iint_S y^2 \mu(x, y, z) dS.$</p>
	 DOWNLOAD DOUBTNUT TODAY FOR FREE PDFs & MORE
288	<p>Moments of Inertia about the x-axis, y-axis, and z-axis $I_x = \iint_S (y^2 + z^2) \mu(x, y, z) dS, I_y = \iint_S (x^2 + z^2) \mu(x, y, z) dS, I_z = \iint_S (x^2 + y^2) \mu(x, y, z) dS,$</p>
289	<p>Volume of a solid Bounded by a closed surface $V = \frac{1}{3} \left \oiint_S xy dz + yz dx + zx dy \right$</p>
290	<p>Gravitational Force $\vec{F} = G \int \int_S \mu(x, y, z) \frac{\vec{r}}{r^3} dS, \text{ where } m \text{ is a mass at a point } (x_0, y_0, z_0) \text{ outside the surface, } \vec{r} = (x - x_0, y - y_0, z - z_0), \mu(x, y, z) \text{ is the density function, and } G \text{ is a gravitational constant.}$</p>
291	<p>Pressure Force $\vec{F} = \iint_S p(\vec{r}) d\vec{S}, \text{ where the pressure } p(\vec{r}) \text{ acts on the surface } S \text{ given by the position vector } \vec{r}.$</p>
292	<p>Fluid Flux (across the surface S) $\Phi = \oiint_S \vec{v}(\vec{r}) \cdot d\vec{S}, \text{ where } \vec{v}(\vec{r}) \text{ is the fluid velocity.}$</p>
293	

Mass Flux (across the surface S) $\Phi = \iint_S \rho \vec{v}(\vec{r}) \cdot d\vec{S}$, where $\vec{F} = \rho \vec{v}$ is the vector field, rho is the fluid density,

294 Surface Charge $Q = \iint_S \sigma(x, y) dS$, where $\sigma(x, y)$ is the surface charge density.

295 Gauss' law the electric flux through any closed surface is proportional to the charge Q enclosed by the surface $\Phi = \iint_S \vec{E} \cdot d\vec{S} = \frac{Q}{\epsilon_0}$ where Φ is the electric flux, \vec{E} is the magnitude of the electric field strength, $\epsilon_0 = 8.85 \times 10^{-12} \frac{F}{m}$ is permittivity of free space.

[Download Hundreds of such PDFs for FREE on Doubtnut App Today](#)

[Download Doubtnut to Ask Any Math Question By just a click](#)

[Get A Video Solution For Free in Seconds](#)

[Doubtnut Has More Than 1 Lakh Video Solutions](#)

[Free Video Solutions of NCERT, RD Sharma, RS Aggarwal, Gengage \(G.Tewani\), Resonance DPP, Allen, Bansal, FIITJEE, Akash, Narayana, VidyaMandir](#)

[Download Doubtnut Today](#)

Get Answer just with a click!

doubtnut has more than 1 Lakh Video Solutions

Update the App now!

GET IT ON Google Play

The advertisement features a pink background with a central yellow box containing the text 'doubtnut has more than 1 Lakh Video Solutions'. To the left and right are two smartphones displaying the app's interface. The left phone shows a math problem and a video solution player. The right phone shows a library of NCERT questions and tips & tricks. At the bottom center is a 'GET IT ON Google Play' button.