

Ques No.

Question

**JEE MAINS 08 APRIL 2019 - PAPER 1 SHIFT 1 -
MEMORY BASED - CHEMISTRY**

Which of the following amine will be prepared by Gabriel phthalimide reaction?

- (1) n-Butylamine
- (2) Triethylamine
- (3) neo-Pentylamine
- (4) tert-Butylamine

CORRECT OPTION: 1

[Watch Free Video Solution on Doubtnut](#)

1 - 9364341

2 - 9364342

**JEE MAINS 08 APRIL 2019 - PAPER 1 SHIFT 1 -
MEMORY BASED - CHEMISTRY**

Reaction of dilute HCl with maltose gives:


- (1) D-glucose
 - (2) D-fructose
 - (3) D-glucose and D-fructose
 - (4) D-galactose
-

CORRECT OPTION: 1

[▶ Watch Free Video Solution on Doubtnut](#)

**JEE MAINS 08 APRIL 2019 - PAPER 1 SHIFT 1 -
MEMORY BASED - CHEMISTRY**

The correct order of K_b value of following is




- (1) $1 > 2 > 3$
 - (2) $1 > 3 > 2$
 - (3) $3 > 2 > 1$
 - (4) $3 > 1 > 2$
-

CORRECT OPTION: 2

[▶ Watch Free Video Solution on Doubtnut](#)

**JEE MAINS 08 APRIL 2019 - PAPER 1 SHIFT 1 -
MEMORY BASED - CHEMISTRY**


Write IUPAC name of the following compound




- (1) 3-hydroxy-2-methylpentanoic acid
- (2) 4-Methyl-3-hydroxypentanoic acid.
- (3) 3-Hydroxy-4-methylpentanoic acid.
- (4) 2-Methyl-2-hydroxypentanoic acid

CORRECT OPTION: 3


[📺 Watch Free Video Solution on Doubtnut](#)





Compound X will be



(1)




(2)




CORRECT OPTION: 2


[▶ Watch Free Video Solution on Doubtnut](#)




X will be:




(1)



(2)





(3)




(4)

CORRECT OPTION: 3


[👁 Watch Free Video Solution on Doubtnut](#)




(1)



(2)



(3) - - -



CORRECT OPTION: 3

[Watch Free Video Solution on Doubtnut](#)

Get Answer just with a click!


doubtNut
has more than
1 Lakh Video
Solutions


Update the App now!

GET IT ON
Google Play



8 - 9364350


**JEE MAINS 08 APRIL 2019 - PAPER 1 SHIFT 1 -
MEMORY BASED - CHEMISTRY**





(1)





CORRECT OPTION: 1

[Watch Free Video Solution on Doubtnut](#)

9 - 9364351


**JEE MAINS 08 APRIL 2019 - PAPER 1 SHIFT 1 -
MEMORY BASED - CHEMISTRY**

Find the compound X which give following test


neutral $\text{FeCl}_3 \rightarrow -ve$

Fehling solution $\rightarrow -ve$



Iodoform reaction $\rightarrow +ve$, brgt Grignard reagent
 $\rightarrow +ve$



(1)




(2)



CORRECT OPTION: 1

[Watch Free Video Solution on Doubtnut](#)

Identify the compound X



- (1) Toluidine
- (2) Benzamide
- (3) Para-cresol
- (4) Oleic acid

CORRECT OPTION: 3

[Watch Free Video Solution on Doubtnut](#)

11 - 9364355

**JEE MAINS 08 APRIL 2019 - PAPER 1 SHIFT 1 -
MEMORY BASED - CHEMISTRY**

Which of the following is not correct for an ideal gas as per first law of thermodynamics

(1) Adiabatic $\Delta U = -w$

(2) Isothermic $q = -w$

(3) cyclic $q = -w$

(4) isobaric $\Delta U = q$


CORRECT OPTION: 1

[▶ Watch Free Video Solution on Doubtnut](#)

12 - 9364357

**JEE MAINS 08 APRIL 2019 - PAPER 1 SHIFT 1 -
MEMORY BASED - CHEMISTRY**

In Freundlich isotherm $\frac{x}{m} \propto P^a$ find the value of a from
the following graph



- (1) $\frac{2}{3}$
- (2) $\frac{1}{3}$
- (3) $\frac{3}{2}$
- (4) 1

CORRECT OPTION: 1

[▶ Watch Free Video Solution on Doubtnut](#)

In a mixture of A and B having vapour pressure of pure A and B are 400 mm Hg and 600 mm Hg respectively. Mole fraction of B in liquid phase is 0.5. Calculate total vapour pressure and mole fraction of A and B in vapour phase-

- (1) 500, 0.4, 0.6
- (2) 500, 0.5, 0.5
- (3) 450, 0.4, 0.6
- (4) 450, 0.5, 0.5

CORRECT OPTION: 1

[Watch Free Video Solution on DoubtNut](#)

14 - 9364360

JEE MAINS 08 APRIL 2019 - PAPER 1 SHIFT 1 - MEMORY BASED - CHEMISTRY

Arrange the following set of quantum number highest energy of an electron.

$$(p) n = 4 \quad i = 1 \quad m = +1 \quad s = +\frac{1}{2}$$

$$(1) n = 4 \quad i = 2 \quad m = -1 \quad s = -\frac{1}{2}$$

$$(r) n = 3 \quad i = 2 \quad m = 0 \quad s = +\frac{1}{2}$$

$$(s) n = 3 \quad i = 1 \quad m = +1 \quad s = -\frac{1}{2}$$

$$(1) q > r > p > s$$

$$(2) q > p > r > s \quad (3) s > p > r > q$$




$$(4) s > r > p > q$$

CORRECT OPTION: 2

[▶ Watch Free Video Solution on Doubtnut](#)


**JEE MAINS 08 APRIL 2019 - PAPER 1 SHIFT 1 -
MEMORY BASED - CHEMISTRY**

A forms ccp lattice B occupy half of the octahedral voids and O occupy all the tetrahedral voids. Calculate formula-



CORRECT OPTION: 1

[▶ Watch Free Video Solution on Doubtnut](#)



**JEE MAINS 08 APRIL 2019 - PAPER 1 SHIFT 1 -
MEMORY BASED - CHEMISTRY**

B_2H_6 reacts with O_2 and H_2O respectively to form

- (1) B_2O_3, H_3BO_3
- (2) B_2O_3, BH_4^-
- (3) HBO_2, H_3BO_3
- (4) H_3BO_3, HBO_2

CORRECT OPTION: 1

[▶ Watch Free Video Solution on DoubtNut](#)

16 - 9364363

17 - 9364365

**JEE MAINS 08 APRIL 2019 - PAPER 1 SHIFT 1 -
MEMORY BASED - CHEMISTRY**

Solution of 100 ml water contains 0.73 g of $Mg(HCO_3)_2$ and 0.81g of $Ca(HCO_3)_2$ calculate the hardness in terms of ppm of $CaCO_3$

(1) $10^2 p \pm$

(2) $10^4 p \pm$

(3) $5 \times 10^3 p \pm$

(4) $10^3 p \pm$

CORRECT OPTION: 2

[📺 Watch Free Video Solution on Doubtnut](#)

18 - 9364366

JEE MAINS 08 APRIL 2019 - PAPER 1 SHIFT 1 - MEMORY BASED - CHEMISTRY

for $Zr_3(PO_4)_4$ the solubility product is K_{sp} and solubility is S. find the correct relation

(1) $S = \left(\frac{K_{sp}}{6912} \right)^{1/7}$

(2) $S = \left(\frac{K_{sp}}{216} \right)^{1/7}$

$$(3) S = \left(\frac{K_{sp}}{216} \right)^{1/8}$$

$$(4) S = \left(\frac{K_{sp}}{912} \right)^{1/3}$$



CORRECT OPTION: 1

[Watch Free Video Solution on Doubnut](#)





19 - 9364367

**JEE MAINS 08 APRIL 2019 - PAPER 1 SHIFT 1 -
MEMORY BASED - CHEMISTRY**

Given complexes are low spin complexes



Then of magnetic (μ) for Fe^{2+} , Cr^{+3} , Ru^{3+} is



CORRECT OPTION: 3

[▶ Watch Free Video Solution on Doubtnut](#)

**JEE MAINS 08 APRIL 2019 - PAPER 1 SHIFT 1 -
MEMORY BASED - CHEMISTRY**

given $E_{S_2O_8^{2-} / SO_4^{2-}}^{\circ} = 2.05V$

$E_{Br_2 / Br^-}^{\circ} = 1.40V$

$E_{Au^{3+} / Au}^{\circ} = 1.10V$, brgt $E_{O_2 / H_2O}^{\circ} = 1.20V$

Which of the following is the strongest oxidizing agent ?

20 - 9364369

(1) $S_2O_8^{2-}$

(2) Br_2

(3) Au^{+3}

(4) O_2

CORRECT OPTION: 1

[▶ Watch Free Video Solution on Doubtnut](#)

21 - 9364370

**JEE MAINS 08 APRIL 2019 - PAPER 1 SHIFT 1 -
MEMORY BASED - CHEMISTRY**

Assertion : Ozone is getting depleted due to CFCs

Reason: With depletion of ozone layer more UV radiation filters into troposphere

(1) both assertion and reason are true and reason is the correct explanation of assertion.

(2) both assertion and reason are true but reason is not correct explanation of assertion.

(3) Assertion is true but reason is false.

(4) Assertion is false but reason is true


CORRECT OPTION: 2

[Watch Free Video Solution on Doubtnut](#)

22 - 9364373

**JEE MAINS 08 APRIL 2019 - PAPER 1 SHIFT 1 -
MEMORY BASED - CHEMISTRY**

In which of the following order of hydrogen energy correct?





CORRECT OPTION: 1

[▶ Watch Free Video Solution on Doubtnut](#)

**JEE MAINS 08 APRIL 2019 - PAPER 1 SHIFT 1 -
MEMORY BASED - CHEMISTRY**

Ellingham diagram is used for

- (1) reduction
- (2) electrolysis
- (3) zone refining
- (4) Van-arkel

CORRECT OPTION: 1

[▶ Watch Free Video Solution on Doubtnut](#)



JEE MAINS 08 APRIL 2019 - PAPER 1 SHIFT 1 - MEMORY BASED - CHEMISTRY

In isoelectronic species Cl^- , Ar , Ca^{2+} size differ due to

- (1) Nuclear charge
- (2) electronic-electronic repulsion in valence shell
- (3) Magnetic quantum number principal number
- (4)

CORRECT OPTION: 1

[Watch Free Video Solution on Doubtnut](#)

24 - 9364375

25 - 9364377

JEE MAINS 08 APRIL 2019 - PAPER 1 SHIFT 1 - MEMORY BASED - CHEMISTRY

3 mole of Ag is heated from 300K to 1000K calculate ΔH when $P = 1 \text{ atm}$ and $C_p = 23 + 0.01T$

(1) $62 \text{ kJ} / \text{mol}$

(2) $45 \text{ kJ} / \text{mol}$

(3) $38 \text{ kJ} / \text{mol}$

(4) $54 \text{ kJ} / \text{mol}$

CORRECT OPTION: 1

[📺 Watch Free Video Solution on Doubtnut](#)

26 - 9364378

**JEE MAINS 08 APRIL 2019 - PAPER 1 SHIFT 1 -
MEMORY BASED - CHEMISTRY**

FeC_2O_4 , $Fe_2(C_2O_4)_3$, $FeSO_4$, $Fe_2(SO_4)_3$ one mole each, will react how many moles of $KMnO_4$.

(1) 1

(2) 2

(3) 3

(4) 5

CORRECT OPTION: 2

[Watch Free Video Solution on Doubtnut](#)

**JEE MAINS 08 APRIL 2019 - PAPER 1 SHIFT 1 -
MEMORY BASED - CHEMISTRY**

$[A]$	$[B]$	Initial Rate
0.05	0.05	0.045
0.10	0.05	0.09
0.20	0.10	0.72

for $2A + B \rightarrow C$. Find the rate law

27 - 9364379

(1) $R = K[A][B]$

(2) $R = k[A][B]^2$

(3) $R = k[A^2][B]$

(4) $R = k[A]^2[B]^2$


CORRECT OPTION: 2

[Watch Free Video Solution on Doubtnut](#)

28 - 9364380

**JEE MAINS 08 APRIL 2019 - PAPER 1 SHIFT 1 -
MEMORY BASED - CHEMISTRY**

Which of the following lanthanoid ions are coloured



CORRECT OPTION: 2

[▶ Watch Free Video Solution on Doubtnut](#)



[▶ Download Doubtnut to Ask Any Math Question By just a click](#)

[▶ Get A Video Solution For Free in Seconds](#)

[▶ Doubtnut Has More Than 1 Lakh Video Solutions](#)

[▶ Free Video Solutions of NCERT, RD Sharma, RS Aggarwal, Cengage \(G.Tewani\), Resonance DPP, Allen, Bansal, FIITJEE, Akash, Narayana, VidyaMandir](#)

[▶ Download Doubtnut Today](#)



Get Answer just
with a click!

doubtnut
has more than
1 Lakh Video
Solutions

Update the App now!

