

Ques No.

Question

**JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

Sum of the series $1 + 2.3 + 3.5 + 4.7 + \dots \dots$ up to 11 terms is

(A) 942

1 - 9469515

(B) 892

(C) 946

(D) 960

CORRECT OPTION: C

[Watch Free Video Solution on Doubtnut](#)

2 - 9469517

**JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

If $P \Rightarrow (q \vee r)$ is false then truth value of p, q, r is respectively

- (A) TTT
- (B) FFF
- (C) FTT
- (D) TFF

CORRECT OPTION: D

[Watch Free Video Solution on DoubtNut](#)

3 - 9469518

**JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

If the ratio of coefficients of the three consecutive terms in binomial expansion of $(1 + x)^n$ is 2:5:70. Then the average of these coefficients is

- (A) 227

(B) 232

(C) 964

(D) 804

CORRECT OPTION: B

[Watch Free Video Solution on DoubtNut](#)

4 - 9469519

JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 -

MEMORY BASED - MATHS

If $\int e^{\sec x} \left(\sec x \tan x f(x) + \sec x \tan x + \tan^2 x \right) dx = e^{\sec x} f(x) + c$

$$\int e^{\sec x} (\sec x \tan x f(x) + \sec x \tan x + \tan^2 x) dx$$

$$= e^{\sec x} f(x) + c$$

. Then $f(x)$ is:

(A) $\sec x + x \tan x + \frac{1}{2}$

(B) $x \sec x + \tan x + \frac{1}{2}$

(C) $x\sec x + x^2\tan x + \frac{1}{2}$

(D) $\sec x + \tan x + \frac{1}{2}$

CORRECT OPTION: D

[Watch Free Video Solution on DoubtNut](#)

5 - 9469520

**JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

The domain of the function

$$f(x) = \frac{1}{x^2 - 4} + n(x^3 - x)$$

(A) $[-1, 0] \cup (1, \infty)$

(B) $\left(-1, \frac{1}{2}\right) \cup (1, 2) \cup (2, \infty)$

(C) $(-1, 0) \cup (1, 2) \cup (2, \infty)$

(D) $(-\infty, 0) \cup (1, 2) \cup (2, \infty)$

CORRECT OPTION: C

[⌚ Watch Free Video Solution on Doubtnut](#)

**JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

The area enclosed by the region $\frac{y^2}{2} \leq x \leq y + 4$ is

- (A) 16
- (B) 8
- (C) 18
- (D) 12

6 - 9469521

CORRECT OPTION: C

[⌚ Watch Free Video Solution on Doubtnut](#)

7 - 9469522

**JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**


Let $A = \begin{bmatrix} 0 & 2y & 1 \\ 2x & y & -1 \\ 2x & -y & 1 \end{bmatrix}$ then number of possible matrices that

$AA = 3I_3$ is

- (A) 3
- (B) 8
- (C) 6
- (D) 4

CORRECT OPTION: D

[⌚ Watch Free Video Solution on Doubtnut](#)



8 - 9469523

JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 - MEMORY BASED - MATHS

The point lying on common tangent to the circles $x^2 + y^2 = 4$

and $x^2 + y^2 + 6x + 8y - 24 = 0$ is

- (A) (6, - 2)
- (B) (4, - 2)
- (C) (4, 6)
- (D) (4)(2, 4)

CORRECT OPTION: A

[⌚ Watch Free Video Solution on Doubtnut](#)

JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 - MEMORY BASED - MATHS

If complex number

$$\omega = \frac{5 + 3z}{5(1 - z)}$$

$|z| < 1$ then

9 - 9469524

- (A) $5Im(\omega) < 1$
- (B) $5Im(\omega) > 4$
- (C) $5Re(\omega) > 1$
- (D) $5Re(\omega) > 1$

CORRECT OPTION: D

[⌚ Watch Free Video Solution on Doubtnut](#)

10 - 9469525

JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 - MEMORY BASED - MATHS

Let α & β are the roots of $(m^2 + 1)x^2 - 3x + (m + 1)^2 = 0$. If sum of roots is maximum then $|\alpha^2 - \beta^3| =$

(A) $8\sqrt{5}$

(B) $8\sqrt{3}$

(C) $10\sqrt{5}$

(D) $4\sqrt{3}$

CORRECT OPTION: A

[Watch Free Video Solution on DoubtNut](#)

11 - 9469526

**JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

If a line makes an angle $\frac{\pi}{3}$ with x -axis, $\frac{\pi}{4}$ with y -axis then angle made by line with z -axis is

- (A) $\frac{2\pi}{3}$
- (B) $(5\pi) \cdot (12)$
- (C) $\frac{3\pi}{4}$
- (D) $\frac{\pi}{12}$

CORRECT OPTION: A

[Watch Free Video Solution on DoubtNut](#)

12 - 9469527

**JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

If two points B and C at distance of 5 units from each other, lie on the line $\frac{x+2}{3} = \frac{y-1}{0} = \frac{2}{4}$. If point A has coordinates $(1, -1, 2)$. Then the area of ΔABC is

- (A) $\sqrt{34}$
- (B) $2\sqrt{34}$

(C) $\sqrt{41}$

(D) $3\sqrt{34}$

CORRECT OPTION: A

[Watch Free Video Solution on DoubtNut](#)

13 - 9469528

JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 - MEMORY BASED - MATHS

If $y(x)$ satisfies the differential equation $\cos x \frac{dy}{dx} - y \sin x = 6x$

. And $y\left(\frac{\pi}{3}\right) = 0$. Then value of $y\left(\frac{\pi}{6}\right)$ is

(A) $\frac{\pi^2}{3\sqrt{2}}$

(B) $\frac{-\pi^2}{2\sqrt{3}}$

(C) $\frac{\pi^2}{2\sqrt{3}}$

(D) $\frac{\pi^2}{4}$

CORRECT OPTION: B

[Watch Free Video Solution on DoubtNut](#)

14 - 9469529

JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 - MEMORY BASED - MATHS

There are two newspaper $A\&B$ published in a city. If 25 % people read A , 20 % people read B & 8 % both read. Also 30 % of these who read A both B look into advertisement 40 % of these who read B but not A look into advertisement 50 % of these who read both $A\&B$ look into advertisement then how many % of people look into advertisement

- (A) 13.9 %
- (B) 13 %
- (C) 13.1 %
- (D) 13.2 %

CORRECT OPTION: A

[⌚ Watch Free Video Solution on Doubtnut](#)

JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 - MEMORY BASED - MATHS

There are two towers of $5m$ & $10m$. The line joining their tops makes 15° with ground then find the distance between them

15 - 9469530
(A) $5(1 - \sqrt{3})$


(B) $10(1 + \sqrt{3})$

(C) $5(2 + \sqrt{3})$

(D) $2(2 + \sqrt{3})$

CORRECT OPTION: C

[⌚ Watch Free Video Solution on Doubtnut](#)



16 - 9469531

JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 - MEMORY BASED - MATHS

If the sum of first 3 terms of an *A.P.* is 33 and their product is 1155. Then the 11th term of the *A.P.* is

- (A) 25
- (B) 25
- (C) 36
- (D) -36

CORRECT OPTION: A

**JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

An inverted cone with semi vertical angle $\theta = \tan^{-1}\left(\frac{1}{2}\right)$ is being filled with water at the rte of $5\text{cm}^3/\text{min}$. Then the rate of change of height of water when height of water is 10cm is

17 - 9469534

- (A) $\frac{2}{3\pi}\text{cm}/\text{min}$
- (B) $\frac{3}{2\pi}\text{cm}/\text{min}$
- (C) $\frac{1}{5\pi}\text{cm}/\text{min}$
- (D) $\frac{2}{5\pi}\text{cm}/\text{min}$

CORRECT OPTION: C

18 - 9469535

**JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

If points $A(-8, d)$ & $B(6, 5)$ lie on a circle C_1 . The line $3y = x + 7$ is a diameter of C_1 . Rectangle, $ABCD$ which is inscribed inside the circle is completed. Then the area of this rectangle is

- (A) 60
- (B) 84
- (C) 42
- (D) 74

CORRECT OPTION: B

[Watch Free Video Solution on DoubtNut](#)

Let $y^2 = 4x$ is a parabola. Then minimum area of a circle touching parabola at $(1, 2)$ as well as x - axis is

(A) $4\pi(3 - 2\sqrt{2})$

(B) $8\pi(3 - 2\sqrt{2})$

(C) $8\pi(4 - 2\sqrt{2})$

(D) $8\pi(3 - 5\sqrt{2})$

CORRECT OPTION: B

[Watch Free Video Solution on Doubtnut](#)

20 - 9469537

**JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

The mean and median of $10, 22, 26, 29, 34, x, 42, 67, 70, y$ (in increasing order) are 42 and 35 respectively then the value

of $\frac{y}{x}$ is

- (A) $\frac{9}{2}$
(B) $\frac{5}{3}$
(C) $\frac{7}{3}$
(D) $\frac{7}{2}$
-

CORRECT OPTION: C

[Watch Free Video Solution on DoubtNut](#)

21 - 9469538

**JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

If tangent of $y^2 = x$ at (α, β) , where $\beta > 0$ is also a tangent of ellipse $x^2 + 2y^2 = 1$ then value of α is

- (A) $\sqrt{2} - 1$
(B) $\sqrt{2} + 1$
(C) $2\sqrt{2 + 1}$
(D) $2\sqrt{2} - 1$

CORRECT OPTION: B

[⌚ Watch Free Video Solution on Doubtnut](#)

**JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

$$\int_0^1 x \cot^{-1} \left(1 - x^2 + x^4 \right) =$$

(A) $\frac{\pi}{2} - \log 2$

22 - 9469539
(B) $\frac{\pi}{2} + \log s\sqrt{2}$

(C) $\frac{\pi}{4} - \log 2$

(D) $\frac{\pi}{4} - \log \sqrt{2}$

CORRECT OPTION: D

[⌚ Watch Free Video Solution on Doubtnut](#)

23 - 9469540

**JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

The value of $\sin 10^\circ \cdot \sin 50^\circ \cdot \sin 70^\circ$ is

(A) $\frac{1}{36}$

(B) $\frac{1}{32}$

(C) $\frac{1}{18}$

(D) $\frac{1}{16}$

CORRECT OPTION: D

[Watch Free Video Solution on Doubtnut](#)

 **doubtnut**
पढ़ना हुआ आसान



Get Solutions as
YOU TYPE

Click Here to
TYPE & ASK

MEMORY BASED - MATHS

$$\text{if } f(x) = \begin{cases} a|\pi - x| + 1 & x \leq 5 \\ b|x - \pi| + 3 & x > 5 \end{cases}$$

(A) $\frac{1}{5 - \pi}$

(B) $\frac{2}{5 - \pi}$

(C) $\frac{2}{5 + \pi}$

(D) $\frac{1}{5 + \pi}$

CORRECT OPTION: B

[Watch Free Video Solution on DoubtNut](#)

25 - 9469542

JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 - MEMORY BASED - MATHS

If the lines $x + (a - 1)y = 1$ and $2x + 1a^2y = 1$ there

$a \in R - \{0, 1\}$ are perpendicular to each other, Then distance
of their point of intersection from the origin is

- (A) $\frac{5}{2}$
- (B) $\frac{2}{\sqrt{5}}$
- (C) $\frac{\sqrt{5}}{2}$
- (D) $\sqrt{2} \frac{2}{5}$

CORRECT OPTION: D

[Watch Free Video Solution on Doubtnut](#)

26 - 9469543

**JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

If some balls are arranged in the form of an equilateral triangle with n rows such that r^{th} row of balls has r balls in it.

If 99 more balls are added to existing one and the balls can now form a square with $(n - 2)$ balls in each row. Then the number of balls that formed the equilateral triangle.

(A) 190

(B) 290

(C) 100

(D) 140

CORRECT OPTION: A

[Watch Free Video Solution on DoubtNut](#)

27 - 9469544

JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 - MEMORY BASED - MATHS

Let $f(2) = 6$ the value of $\lim_{x \rightarrow 2} \int_6^{f(x)} \frac{2tdt}{x-2}$

(A) $12f'(2)$

(B) $24f'(2)$

(C) $8f'(2)$

(D) $10f'(2)$

CORRECT OPTION: A

[⌚ Watch Free Video Solution on Doubtnut](#)

JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 - MEMORY BASED - MATHS

if $f(x) = [x] - \left[\frac{x}{4} \right]$ then

- 28 - 9469545
- (A) $f(x)$ exists and $\lim_{x \rightarrow 4} f(x)$ both does not exist
 - (B) function $f(x)$ is continuous at $x = 4$
 - (C) $f(x)$ exists and $\lim_{x \rightarrow 4^-} f(x)$ both exist but are not equal
 - (D) $f(x)$ exists but $\lim_{x \rightarrow 4^-} f(x)$ does not exist

CORRECT OPTION: B

[⌚ Watch Free Video Solution on Doubtnut](#)

29 - 9469546

JEE MAINS 09 APRIL 2019 - PAPER 1 SHIFT 2 -

MEMORY BASED - MATHS

Let the system of linear equations

$2x + 3y - z = 0$, $2x + ky - 3z = 0$ and $2x - y + z = 0$ have non

trivial solution then $\frac{x}{y} + \frac{y}{z} + \frac{z}{x} + k$ will be

(A) 2

(B) 3

(C) 1

(D) -4

CORRECT OPTION: B

[Watch Free Video Solution on Doubtnut](#)



- ☛ Download Doubtnut to Ask Any Math Question By just a click
- ☛ Get A Video Solution For Free in Seconds
- ☛ Doubtnut Has More Than 1 Lakh Video Solutions
- ☛ Free Video Solutions of NCERT, RD Sharma, RS Aggarwal, Cengage (G.Tewani), Resonance DPP, Allen, Bansal, FIITJEE, Akash, Narayana, Vidyamandir
- ☛ Download Doubtnut Today

Get Answer just
with a click!

doubtnut
has more than
1 Lakh Video
Solutions

Update the App now!

GET IT ON
Google Play

