

Ques No.

Question

**JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

The locus of centre of circle which touches the circle

$x^2 + y^2 = 1$ and y – axis in 1st quadrant is

(A) $y^2 = 2x - 1$

(B) $x^2 + 1 = 2y$

(C) $y^2 = 2x + 1$

(D) $x^2 - 1 = 2y$

1 - 9525896

CORRECT OPTION: C

[Watch Free Video Solution on Doubtnut](#)

2 - 9525897

JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 -

MEMORY BASED - MATHS

Negation of statement $-s \vee (-r \wedge s)$ is

- (A) $s \wedge r$
- (B) $s \vee r$
- (C) $-s \rightarrow r$
- (D) $-s \wedge r$

CORRECT OPTION: A

[⌚ Watch Free Video Solution on DoubtNut](#)

3 - 9525898

JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 - MEMORY BASED - MATHS

There are 20 pillars of equal height on a circular ground any two non adjacent pillars joined by a beam. Then the number of such beams are:

- (A) 180
(B) 210
(C) $.^{20} C_2 - 20$
(D) $.^{20} C_2$
-

CORRECT OPTION: C

[Watch Free Video Solution on Doubtnut](#)

4 - 9525899

**JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

If one of the directic of hyperbola $\frac{x^2}{9} - \frac{y^2}{16} = 1$ is $x = -\frac{9}{5}$ then the corresponding of hyperbola is

- (A) (5, 0)
(B) (-5, 0)
(C) (0, 4)
(D) (0, -4)

CORRECT OPTION: B

[⌚ Watch Free Video Solution on Doubtnut](#)

JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 - MEMORY BASED - MATHS

Let Z and w be two complex numbers such that $|zw| = 1$ and $\arg(z) - \arg(w) = \pi/2$ then

(A) $z\bar{w} = -i$

5 - 9525900
(B) $z\bar{w} = i$

(C) $z\bar{w} = \frac{1-i}{\sqrt{2}}$

(D) $z\bar{w} = \frac{1+i}{\sqrt{2}}$

CORRECT OPTION: B

[⌚ Watch Free Video Solution on Doubtnut](#)

JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS

A straight line parallel to the straight line $4x - 3y + 2 = 0$ is at a distance $\frac{3}{5}$ units from the origin. Then which of the following point lie on this line.

- (A) $\left(\frac{1}{4}, \frac{2}{3}\right)$
- (B) $\left(-\frac{1}{4}, \frac{2}{3}\right)$
- (C) $\left(\frac{1}{4}, -\frac{1}{3}\right)$
- (D) $\left(-\frac{1}{4}, -\frac{2}{3}\right)$

CORRECT OPTION: B

[⌚ Watch Free Video Solution on DoubtNut](#)

Value of $\int_{\pi/6}^{\pi/3} \sec^{2/3} x \cdot \csc^{4/3} x dx$ is

(A) $2^{\frac{7}{6}} - 2^{\frac{5}{6}}$

(B) $2^{\frac{5}{6}} - 2^{\frac{3}{4}}$


(C) $3^{\frac{5}{6}} - 3^{\frac{3}{4}}$

(D) $3^{\frac{7}{6}} - 3^{\frac{5}{6}}$

CORRECT OPTION: D

[Watch Free Video Solution on DoubtNut](#)

 doubtNut
पढ़ना हुआ आसान



8 - 9525903

**JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

If $\lim_{x \rightarrow 1} \frac{x^2 - ax + b}{x - 1} = 5$ then the value of $a + b$ is

- (A) 1
- (B) -5
- (C) 4
- (D) -7

CORRECT OPTION: D

[Watch Free Video Solution on DoubtNut](#)

9 - 9525904

**JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

If both the standard deviation and mean of data $x_1, x_2, x_3, \dots, x_{50}$ are 16, then the mean of the data set $(x_1 - 4)^2, (x_2 - 4)^2, (x_3 - 4)^2, \dots, (x_{50} - 4)^2$ is

- (A) 200
 - (B) 100
 - (C) 400
 - (D) 1600
-

CORRECT OPTION: C

[Watch Free Video Solution on Doubtnut](#)

10 - 9525905

**JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

A coin is tossed n times, if the probability of getting at least one head is atleast 99 % then the minimum value of n is

- (A) 6
- (B) 7
- (C) 8
- (D) 9

CORRECT OPTION: B

[⌚ Watch Free Video Solution on Doubtnut](#)

**JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

if $\int x^5 e^{-x^2} + C$ then the value of $g(-1)$ is

- 11 - 9525906
- (A) $\frac{3}{2}$
 - (B) $\frac{1}{2}$
 - (C) $-\frac{5}{2}$
 - (D) $\frac{e}{2}$

CORRECT OPTION: C

[⌚ Watch Free Video Solution on Doubtnut](#)

12 - 9525907

**JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

The solution differential equation

$\frac{dy}{dx} + y \tan x = 2x + x^2 \tan x$ is

(A) $y = x^2 + c \cos x$

(B) $y = 2x^2 - c \cos x$

(C) $y + x^2 = c \cos x$

(D) $y + 2x^2 = c \cos x$

CORRECT OPTION: A

[Watch Free Video Solution on DoubtNut](#)

13 - 9525908

**JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

Let common tangent to curves $x^2 + y^2 = 1$ and

$y^2 = 4\sqrt{2}x$ is $y = -ax + c$ then $|c|$ is equal

(A) $\sqrt{2}$

(B) 1

(C) $\frac{1}{\sqrt{2}}$

(D) 2

CORRECT OPTION: A

[Watch Free Video Solution on Doubtnut](#)

14 - 9525909

**JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

Area of triangle formed by tangent and normal to ellipse

$3x^2 + 5y^2 = 32$ at point (2, 2) and x-axis is

(A) $\frac{68}{15}$

(B) $\frac{36}{15}$

(C) $\frac{32}{3}$

(D) $\frac{31}{3}$

CORRECT OPTION: A

[⌚ Watch Free Video Solution on Doubtnut](#)


**JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

Area enclosed by curves $y = 2^x$ and $y = |x + 1|$ in the first quadrant is

- 15 - 9525910
- (A) $\frac{1}{2} - \frac{1}{\log_2}$
 - (B) $\frac{3}{2} - \frac{1}{2 \log 2}$
 - (C) $\frac{3}{2} - \frac{1}{\log 2}$
 - (D) $\frac{1}{2} + \frac{3}{\log 2}$

CORRECT OPTION: C

[⌚ Watch Free Video Solution on Doubtnut](#)



16 - 9525911

JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 - MEMORY BASED - MATHS

if the foot of perpendicular drawn from a point on the line

$$\frac{x-1}{2} = \frac{y+1}{-1} = \frac{z}{1}$$

on the plane $x + y + z = 3$ also lies on the plane $x - y + z = 3$ then the coordinates of the foot of perpendicular is

(A) (-2, 0, 5)

(B) (-1, 0, 4)

(C) (1, 0, 2)

(D) (2, 0, 1)

CORRECT OPTION: D

[Watch Free Video Solution on DoubtNut](#)

17 - 9525912

JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 - MEMORY BASED - MATHS

If three parallel planes are given by

$$P_1: 2x - y + 2z = 6$$

$$P_2: 4x - 2y + 4z = \lambda$$

$$P_3: 2x - y + 2z = \mu$$

if distance between P_1 and P_2 is $\frac{1}{3}$ and between P_1 and

P_3 is $\frac{2}{3}$ then the maximum value of $\lambda + \mu$ is

(A) 22

(B) 20

(C) 18

(D) 24

CORRECT OPTION: A

[Watch Free Video Solution on Doubtnut](#)

JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 - MEMORY BASED - MATHS

If terms $a_1, a_2, a_3, \dots, a_{50}$ are in A.P and $a_6 = 2$, then the value of common difference at which maximum value of $a_1 a_4 a_5$ occur is

- 18 - 9525913
- (A) $\frac{3}{5}$
 - (B) $\frac{8}{5}$
 - (C) $\frac{2}{5}$
 - (D) $\frac{2}{3}$

CORRECT OPTION: B

[Watch Free Video Solution on Doubtnut](#)

19 - 9525914

JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 -

MEMORY BASED - MATHS

if a, b, c are in G.P. and $3a, 7b, 15c$ are first 3 term of A.P.

Also the common ratio of G.P. $\in \left(0, \frac{1}{2}\right)$ then the 4th term of A.P is

- (A) $\frac{3}{2}a$
- (B) $\frac{7}{2}a$
- (C) $\frac{5}{2}a$
- (D) a

CORRECT OPTION: D

[Watch Free Video Solution on DoubtNut](#)

In a $\triangle ABC$, $c = 4$ and angles A, B and C are in A.P
also ratio $a : b$ is $1 : \sqrt{3}$ then area of $\triangle ABC$ is

(A) $\sqrt{3}$

(B) $2\sqrt{3}$

(C) $3\sqrt{3}$

(D) $4\sqrt{3}$

CORRECT OPTION: B

[Watch Free Video Solution on DoubtNut](#)

21 - 9525916

**JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

If the coefficient of x in binomial expansion of expression

$$\left(x^2 + \frac{1}{x^3}\right)^n$$
 is $.^n C_{23}$. Then the minimum value of n is

(A) 28

(B) 48

(C) 58

(D) 38

CORRECT OPTION: D

[⌚ Watch Free Video Solution on DoubtNut](#)

22 - 9525917

JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 -

MEMORY BASED - MATHS

Number of real solution of the equation

$x^2 + |2^n - 1| = 2^x(2^x - 2)$ is

(A) 0

(B) 1

(C) 2

(D) 3

CORRECT OPTION: B

[Watch Free Video Solution on Doubtnut](#)

JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 - MEMORY BASED - MATHS

If $f(x) = \ln(\sin x)$ and $g(x) = \sin^{-1}(e^{-x})$ for all $x \in (0, \pi)$ and $f(g(\alpha)) = b$ and $(f(g(x)))'$ at $x = \alpha$ is a then which is true.

23 - 9525918

- (A) $a\alpha^2 - b\alpha + 1 = a$
- (B) $a\alpha^2 - b\alpha^2 = -a$
- (C) $a\alpha^2 - b\alpha + 1 = -a$
- (D) $a\alpha^2 - b\alpha - 1 = -a$

CORRECT OPTION: C

[Watch Free Video Solution on Doubtnut](#)



24 - 9525919

JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 - MEMORY BASED - MATHS

if

$$\cos^{-1} x - \cos^{-1}(y/2) = \alpha, x \in [-1, 1], y \in [-2, 2]$$

then the value of $4x^2 - 4xy \cos \alpha + y^2$ is

(A) $4 \sin^2 \alpha$

(B) $2 \sin^2 \alpha$

(C) $4 \cos^2 \alpha$

(D) $2 \cos^2 \cos^2 \alpha$

CORRECT OPTION: A

[Watch Free Video Solution on Doubtnut](#)

JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 - MEMORY BASED - MATHS

A spherical ball of radius 10 cm is enclosed by ice of uniform thickness in spherical shape. If ice melts at the rate of $50\text{cm}^3 / \text{min}$ then the rate of decrease of thickness of ice when thickness of ice is 5 cm is

25 - 9525920

- (A) $\frac{1}{36\pi}\text{cm} / \text{min}$
- (B) $\frac{1}{9\pi}\text{cm} / \text{min}$
- (C) $\frac{1}{18\pi}\text{cm} / \text{min}$
- (D) $\frac{2}{9\pi}\text{cm} / \text{min}$

CORRECT OPTION: C

[Watch Free Video Solution on Doubtnut](#)

**JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

If a tangent is drawn parallel to the line $6x + 18y - 11 = 0$ to the curve $y = \frac{x}{x^2 - 3}$ which touches the curve at point (α, β) then

- (A) $|6\alpha + 2\beta| = 19$
- (B) $|6\alpha + 2\beta| = 11$
- (C) $|2\alpha + 6\beta| = 7$
- (D) $|2\alpha + 6\beta| = 11$

CORRECT OPTION: A

[⌚ Watch Free Video Solution on DoubtNut](#)

**JEE MAINS 10 APRIL 2019 - PAPER 1 SHIFT 2 -
MEMORY BASED - MATHS**

value of

$$1 + \frac{1^3 + 2^3}{1+2} + \frac{1^3 + 2^3 + 3^3}{1+2+3} + \dots \dots$$

$$+ \frac{1^3 + 2^3 + \dots + 15^3}{1+2+\dots+15} - \frac{1}{2}(1+2+\dots+15)$$

is

(A) 840

(B) 720

(C) 540

(D) 680

CORRECT OPTION: D

[Watch Free Video Solution on Doubtnut](#)



- 👉 Download Doubtnut to Ask Any Math Question By just a click
- 👉 Get A Video Solution For Free in Seconds
- 👉 Doubtnut Has More Than 1 Lakh Video Solutions
- 👉 Free Video Solutions of NCERT, RD Sharma, RS Aggarwal, Cengage (G.Tewani), Resonance DPP, Allen, Bansal, FIITJEE, Akash, Narayana, Vidyamandir
- 📲 Download Doubtnut Today

Get Answer just
with a click!

doubtnut
has more than
1 Lakh Video
Solutions

Update the App now!

GET IT ON
Google Play

